



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 265/26, C07C 271/28, A61K 31/536, A61P 3/04</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/40569</b> <b>(43) International Publication Date:</b> 13 July 2000 (13.07.00)
<b>(21) International Application Number:</b> PCT/GB00/00031 <b>(22) International Filing Date:</b> 6 January 2000 (06.01.00) <b>(30) Priority Data:</b> 9900416.0 8 January 1999 (08.01.99) GB <b>(71) Applicant (for all designated States except US):</b> ALIZYME THERAPEUTICS LIMITED [GB/GB]; 280 Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> HODSON, Harold, Francis [GB/GB]; 69 Whitecroft Way, Beckenham, Kent BR3 3AW (GB). DOWNHAM, Robert [GB/GB]; Cambridge Discovery Chemistry, The Merrifield Centre, Rosemary Lane, Cambridge CB1 3LQ (GB). MITCHELL, Timothy, John [GB/GB]; Cambridge Discovery Chemistry, The Merrifield Centre, Rosemary Lane, Cambridge CB1 3LQ (GB). CARR, Beverley, Jane [GB/GB]; Boys Bridge Cottage, Fen Road, Abbingdon Piggotts, Cambs. SG8 5PQ (GB). DUNK, Christopher, Robert [GB/GB]; Alizyme Therapeutics Limited, 280 Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). PALMER, Richard, Michael, John [GB/GB]; Alizyme Therapeutics Limited, 280 Cam-		bridge Science Park, Milton Road, Cambridge CB4 4WE (GB). <b>(74) Agents:</b> FLORENCE, Julia, Anne et al.; Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB). <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>With international search report.</i>

U.S. Appl. No. 10/617,498  
 Filed July 11, 2003; Schoenfinger, et al  
 File: DEAV2002/0050US NP

**(54) Title:** 2-OXY-4H-3,1-BENZOXAZIN-4-ONES FOR TREATMENT OF OBESITY

**(57) Abstract**

The use of a compound comprising formula (I) or a salt, ester, amide or prodrug thereof in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality, e.g. in the control and inhibition of unwanted enzymes in products and processes. The compounds



are also useful in medicine e.g. in the treatment of obesity and related conditions. The invention also relates to novel compounds within formula (I), to processes for preparing them and pharmaceutical compositions containing them. In formula (I) A is a 6-membered aromatic or heteroaromatic ring; and R<sup>1</sup> is a branched or unbranched alkyl (optionally interrupted by one or more oxygen atoms), alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, reduced arylalkyl, arylalkenyl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, reduced aryl, reduced heteroaryl, reduced heteroarylalkyl or a substituted derivative of any of the foregoing groups.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## 2-OXY-4H-3,1-BENZOXAZIN-4-ONES FOR TREATMENT OF OBESITY


The present invention provides known and novel compounds, their use in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality (*in vivo*, as the enzyme naturally occurs) their use in  
5 medicine, and particularly in the prevention and/or treatment of obesity or an obesity-related disorder. Also provided are methods for the prevention and/or treatment of obesity or an obesity-related disorder and for promoting/aiding non-medical weight loss and the use of the compounds in the manufacture of a medicament for the  
10 aforementioned indications. In respect of novel compounds the invention also provides processes for their manufacture, compositions containing them and methods for manufacturing such compositions.

In the last 20 years, there has been an increasing trend in obesity in the populations of  
15 the developed world. The increased incidence of obesity is due in part to the ready availability of food in numerous retail outlets and westernised diets that have high saturated fat and lower fibre contents such that the food is energy dense. The lifestyle of the populations of the developed world has also become more sedentary with the increased mechanisation of society and the steady reduction of manual labour  
20 intensive industries. There now exists an energy imbalance between the energy intake from calorie dense foods and the reduced energy expenditure required for a sedentary lifestyle. Some of the excess energy intake is stored as fat in the adipose tissue, the accumulation of which over a period of time results in obesity and can be a significant contributory factor to other disease and disorders.


25 Obesity is now recognised by the medical profession as a metabolic disease. In the USA, it is estimated that 25% of the adult population is considered clinically obese (Body Mass Index>30). Obesity can be a debilitating condition which reduces the quality of life and increases the risk of related disorders such as diabetes,  
30 cardiovascular disease and hypertension. It has been estimated that \$45 billion of US

healthcare costs, or 8% per annum of total healthcare spend, is as a direct result of obesity. The traditional approaches to long term weight management such as diet and exercise have proved ineffective alone to control the spread of obesity. Today, more than ever, there is considerable interest in developing safe, effective drugs for the treatment of obesity.

Pharmacological approaches to the treatment of obesity have focused on either developing drugs that increase energy expenditure or drugs that reduce energy intake. One approach to the reduction of energy intake is to reduce the body's ability to digest and absorb food, in particular fat. The key enzymes involved in the digestion of fat are hydrolytic enzymes. The most significant of the fat degrading enzymes are lipases, primarily, but not exclusively pancreatic lipase that is secreted by the pancreas into the gut lumen. The lipase inhibitor lipstatin has formed the basis of the anti-obesity drug, orlistat. Orlistat is the subject of published European Patent Application No. EP129748, which relates to compounds of formula:



where A is  $-(CH_2)_5-$  or;




and their use in inhibiting pancreatic lipase and treating hyperlipaemia and obesity. Orlistat has as its major active moiety a beta-lactone group that reacts to form an ester

with the side chain hydroxyl group of serine 152 within the active site of pancreatic lipase.

Even if orlistat provides an effective method for treating obesity, there remains a need to provide alternative drugs and methods for use in the control and treatment of obesity and obesity-related disorders and in promoting or aiding non-medical weight loss. Inhibitors of enzymes involved in the degradation of fat are provided here and shown to be effective in the prevention and/or treatment of obesity, obesity-related disease and/or in promoting cosmetic weight loss.


US Patent No. 4,665,070 (Syntex) describes a broad class of 2-oxy-4*H*-3,1-benzoxazin-4-ones of the formula:



where *a* is an integer from 0-4; each *R'* may be selected from a wide range of substituents; *A* is a bond or a C<sub>1-8</sub> alkylene group; and *R* is H (except when *A* is a bond) phenyl, imidazolyl or C<sub>3-6</sub> cycloalkyl each of which rings may be optionally substituted. The *R'* groups are said to be preferably in the 5- and/or 7-positions of the ring. A preferred value of the group *A* is lower alkylene having 1-4 carbon atoms. In the most preferred compounds *A* is ethylene. The compounds are said to be useful as serine protease inhibitors and to treat physiologic conditions and disease states known to involve enzymes, or as contraceptives. The specification describes various conditions and diseases involving enzymatic pathways, including inflammation, arthritis, tumor cell metastasis, pulmonary emphysema, mucocutaneous lymph node syndrome, adult respiratory distress syndrome and pancreatitis. It is also suggested

that the compounds may have antiparasitic, anticoagulant and/or antiviral activity. Similar compounds are also described in US Patent No. 4,745,116.


- 5 International Patent Application No. WO89/07639 (BP Chemicals Ltd) describes detergent compositions in aqueous solution which comprise a surfactant, a precursor compound capable of giving rise to a peroxygen compound in the presence of water, a suds suppressing agent, a detergent builder and a bleach activator which may have the formula:



- 10 where R is *inter alia* an alkoxy group and R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> (which may be the same or different) are selected from H, halogen, alkyl, alkenyl, aryl, hydroxyl, alkoxy, amino, alkylamino, -COOR<sub>5</sub> and carbonyl functions. The number of carbon atoms in the alkyl groups and moieties is not defined, but the specific examples are of lower alkyl and alkoxy groups e.g. R may be ethoxy.

15


East German Patent No. DD 246996A1 describes a process for preparing 2-alkoxy- and 2-aryloxy-3,1-benzoxazin-4-ones of the formula:



where  $R'_n$  represents one or more H atoms and/or other substituents such as alkyl, alkoxy, aralkyl, aryl, thiocyanato, mercapto, alkylthio, halogen or nitro, and  $R^2$  represents an alkyl, aralkyl or aryl residue. The compounds are said to be useful as herbicides and fungicides and to have activity as inhibitors of chymotrypsin. Specific examples of  $R^2$  are ethyl, benzyl and phenyl.

We have now found that a particular class of benzoxazinone compounds has activity as lipase inhibitors.

Accordingly, in a first aspect, the present invention provides the use of a compound comprising formula (I):



(I)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof; in the manufacture of a medicament for the treatment of conditions which require the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality,

wherein in formula (I):

A is a 6-membered aromatic or heteroaromatic ring; and

$R^1$  is a branched or unbranched alkyl (optionally interrupted by one or more oxygen atoms), alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, reduced arylalkyl, arylalkenyl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, reduced aryl, reduced heteroaryl, reduced heteroarylalkyl or a substituted derivative of any of the foregoing groups, wherein the substituents are one or more independently of halogen, alkyl, halosubstituted alkyl, aryl, arylalkyl, heteroaryl, reduced heteroaryl, reduced heteroarylalkyl, arylalkoxy, cyano, nitro,  $-C(O)R^4$ ,  $-CO_2R^5$ ,  $-SOR^4$ ,  $-SO_2R^4$ ,  $-NR^6R^7$ ,  $-OR^6$ ,  $-SR^6$ ,  $-C(O)CX^1X^2NR^6R^7$ ,  $-C(O)N(OH)R^6$ ,  $-C(O)NR^5R^4$ ,  $-NR^6C(O)R^4$ ,  $-CR^6(NH_2)CO_2R^6$ ,  $-NHCX^1X^2CO_2R^6$ ,  $-N(OH)C(O)NR^6R^7$ ,  $-N(OH)C(O)R^4$ ,  $-NHC(O)NR^6R^7$ ,  $-C(O)NHNHNR^6R^7$ ,  $-C(O)N(OR^5)R^6$ , or a lipid or steroid (natural or synthetic) with the proviso that any hetero atom substituent in  $R^1$  must be separated from the exocyclic oxygen atom by at least two carbon atoms (preferably saturated);

and where:-

$R^4$  is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, reduced heteroaryl, reduced heteroarylalkyl,  $-OR^6$ ,  $-NHCX^1X^2CO_2R^6$  or  $-NR^6R^7$ ;

$R^5$  is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, reduced heteroaryl or reduced heteroarylalkyl; and

$R^6$  and  $R^7$  are independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, reduced heteroaryl, reduced heteroarylalkyl or  $-(CH_2)_n(OR^5)_m$  wherein  $n$  is 1 to 12, preferably 2 to 10, wherein  $m$  is 1-3 and  $R^5$  is most preferably  $C_2-C_{10}$  alkyl; and

$X^1$  and  $X^2$  are independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, reduced heteroaryl or reduced heteroarylalkyl.



In compounds of formula (I) any alkyl, alkenyl and alkynyl groups and moieties may be straight chain (unbranched) or branched chain. Straight chain alkyl, alkenyl and alkynyl groups or moieties may contain from 1 to 30 carbon atoms, eg. 1 to 25 carbon atoms, preferably 1 to 20 carbon atoms. Branched chain alkyl, alkenyl and alkynyl groups or moieties may contain from 1 to 50 carbon atoms, preferably 1 to 30 carbon atoms.

Preferred values for  $R^1$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$ ,  $X^1$  and  $X^2$  are as defined below for formulae (II) and (IIa). In particular, preferred values for  $R^4$ ,  $R^5$  and  $R^6$  are as defined for  $R^{13}$  hereinbelow and preferred values for  $R^7$  are as defined for  $R^{14}$  hereinbelow.

In this text, 'reduced', in the context of 'reduced heteroaryl' and the like means fully or partially saturated.

Aryl groups include for example optionally substituted unsaturated monocyclic or bicyclic rings of up to 12 carbon atoms, such as phenyl and naphthyl, and partially saturated bicyclic rings such as tetrahydro-naphthyl. Examples of substituents which may be present on an aryl group include one or more of halogen, amino, nitro, alkyl, haloalkyl, alkoxy, phenoxy and phenoxy substituted by one or more of halo, alkyl or alkoxy.


A heteroaryl group or moiety may be for example an optionally substituted 5- or 6-membered heterocyclic aromatic ring which may contain from 1 to 4 heteroatoms selected from O, N and S. The heterocyclic ring may optionally be fused to a phenyl ring. Examples of heteroaryl groups thus include furyl, thienyl, pyrrolyl, oxazolyl, oxazinyl, thiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, pyridyl, triazolyl, triazinyl, pyridazyl, pyrimidinyl, pyrazolyl, indolyl, indazolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzoxazinyl, quinoxalyl, quinolyl, quinazolyl, cinnolyl, benzothiazolyl, pyridopyrrolyl. Suitable substituents include one or more

of halogen, oxo, amino, nitro, alkyl, haloalkyl, alkoxy, phenoxy and phenoxy substituted by one or more of halo, alkyl, haloalkyl or alkoxy.

5 A reduced heteroaryl group or moiety may be for example a fully or partially saturated derivative of the aforementioned heteroaryl groups. Examples of reduced heteroaryl groups thus include pyrrolidinyl, tetrahydrofuryl, tetrahydrothienyl and piperidinyl.

10 The compounds of formula (I) are useful inhibitors of enzymes involved in the degradation of fats. Preferably therefore the first aspect of the invention provides the use of a compound of formula (I) as defined hereinabove, or a pharmaceutically acceptable salt, ester, amide or prodrug thereof, in the manufacture of a medicament for the control or treatment of obesity, or obesity-related disorders or for promoting non-medical weight loss.

15 Preferably, a compound for use according to the first aspect of the invention is a compound of formula (II):



20

(II)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof,

wherein:-

$R^1, R^4, R^5, R^6, R^7, X^1$  and  $X^2$  are as defined above for formula (I); and

$R^8, R^9, R^{10}, R^{11}$  are each independently hydrogen, halo, hydroxy, amino, nitro, cyano,

5

or a group  $R^1$ , as defined above,

or a group  $R^{12}Q$  where  $Q$  is O, CO, CONH, NHCO, S, SO,  $SO_2$ , or  $SO_2NH_2$  and  $R^{12}$  is hydrogen or a group  $R^1$  as defined above,

10

or a group  $R^1R^2N$  where  $R^1$  is as defined above and  $R^2$  is hydrogen or  $R^1$ , with the proviso that any hetero atom substituent in  $R^1$  and/or  $R^2$  must be separated from the aromatic hetero atom substituent by at least two carbon atoms (preferably saturated). Preferred values of  $R^1, R^8, R^9, R^{10}$  and  $R^{11}$  are as defined below for compounds of

15

formula (IIa).

More preferably, a compound for use according to the first aspect of the invention comprises a compound of formula (II), or a pharmaceutically acceptable salt, ester, amide or prodrug thereof; wherein:

20

$R^1$  is either a branched or unbranched alkyl group having up to 25, e.g. up to 20 carbon atoms, an aryl (e.g. optionally substituted phenyl or 2-naphthyl), an arylalkyl group wherein the alkyl moiety has up to 25, e.g. up to 20 carbon atoms, or an aryl aryl group, wherein the aryl alkyl group or the aryl aryl group may be separated by a spacer, and where the spacer can be one or more of an ester, amide, O,  $CH_2$  or a ketone and wherein any aryl group is preferably a phenyl, optionally substituted with alkyl, haloalkyl or halogen;

25

$R^8$  is hydrogen or fluorine;

30


$R^9$  is lower branched or unbranched alkyl having 1 to 10 carbon atoms, preferably methyl; cyclic alkyl having 3 to 10 carbon atoms, preferably cyclopropyl; haloalkyl, preferably trifluoromethyl; or a halogen, most preferably chlorine or fluorine;

5  $R^{10}$  is hydrogen lower branched or unbranched alkyl having 1 to 10 carbon atoms, preferably methyl; cyclic alkyl having 3 to 10 carbon atoms, preferably cyclopropyl; haloalkyl, preferably trifluoromethyl; or a halogen, most preferably chlorine or fluorine;

10  $R^{11}$  is hydrogen lower branched or unbranched alkyl having 1 to 10 carbon atoms, preferably methyl, or halogen, preferably fluorine.

In particular,  $R^1$  is an unbranched alkyl group, having 12, 14, 15, 16, 17 or 18 carbon atoms in the alkyl chain. In addition to this particular option for  $R^1$ ,  $R^9$  may be  
15 methyl.

In a second aspect the present invention provides novel compounds of formula (IIa):



(IIa)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof, where:

20  $R^{1a}$  is

- (i) a  $C_{10-30}$  branched or unbranched alkyl, optionally substituted by one or more independently of  $C_{3-6}$  cycloalkyl,  $C_{3-6}$  cycloalkenyl, aryl, heteroaryl, reduced

- heteroaryl,  $-C(O)R^{13}$ ,  $-CO_2R^{13}$ ,  $-SOR^{13}$ ,  $-SO_2R^{13}$ ,  $-NR^{13}R^{14}$ ,  $-OR^{13}$ ,  $-SR^{13}$ ,  $-C(O)NR^{13}R^{14}$ ,  $-NR^{14}C(O)R^{13}$ , halogen, cyano, and nitro and/or optionally interrupted by one or more oxygen atoms with the proviso that any hetero atom in  $R^1$  must be separated from the exocyclic oxygen atom (or from any other heteroatom) by at least two carbon atoms (preferably saturated);
- (ii)  $C_{2-25}$  alkenyl,  $C_{2-25}$  alkynyl,  $C_{3-6}$  cycloalkenyl, aryl- $C_{2-25}$  alkenyl, heteroaryl- $C_{2-25}$  alkenyl, reduced heteroaryl, reduced heteroaryl- $C_{1-25}$  alkyl or a substituted derivative of any of the foregoing groups wherein the substituents are one or more independently of  $C_{1-6}$  alkyl, halosubstituted  $C_{1-6}$  alkyl, aryl, aryl- $C_{1-6}$  alkyl, heteroaryl, reduced heteroaryl, reduced heteroaryl- $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, aryl- $C_{1-6}$  alkoxy,  $-C(O)R^{13}$ ,  $-CO_2R^{13}$ ,  $-SOR^{13}$ ,  $-SO_2R^{13}$ ,  $-NR^{13}R^{14}$ ,  $-OR^{13}$ ,  $-SR^{13}$ ,  $-C(O)NR^{13}R^{14}$ ,  $-NR^{14}C(O)R^{13}$ , halogen, cyano, and nitro, with the proviso that any hetero atom in  $R^1$  must be separated from the exocyclic oxygen atom (or from any other heteroatom) by at least two carbon atoms (preferably saturated);
- (iii) a  $C_{1-9}$  alkyl group interrupted by one or more oxygen atoms and optionally substituted by one or more independently of  $C_{3-6}$  cycloalkyl,  $C_{3-6}$  cycloalkenyl, aryl, heteroaryl, reduced heteroaryl,  $-C(O)R^{13}$ ,  $-CO_2R^{13}$ ,  $-SOR^{13}$ ,  $-SO_2R^{13}$ ,  $NR^{13}R^{14}$ ,  $OR^{13}$ ,  $SR^{13}$ ,  $-C(O)NR^{13}R^{14}$ ,  $-NR^{14}C(O)R^{13}$ , halogen, cyano and nitro with the proviso that any hetero atom in  $R^1$  must be separated from the exocyclic oxygen atom (or from any other heteroatom) by at least two carbon atoms (preferably saturated); or
- (iv) a  $C_{1-9}$  alkyl group substituted by a group selected from  $-C(O)R^{13}$ ,  $-CO_2R^{13}$ ,  $SOR^{13}$ ,  $SO_2R^{13}$ ,  $NR^{13}R^{14}$ ,  $OR^{13}$ ,  $SR^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ; halogen, cyano, nitro, bicyclic aryl, bicyclic heteroaryl, monocyclic or bicyclic reduced heteroaryl, and monocyclic heteroaryl other than imidazolyl;

(v) a phenyl group substituted by a group selected from  $\text{OR}^{17}$ ,  $-\text{COR}^{13}$ ,  $-\text{CO}_2\text{R}^{13}$ ,  $\text{SOR}^{13}$ ,  $\text{SO}_2\text{R}^{13}$ ,  $\text{CONR}^{13}\text{R}^{14}$ ,  $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$ ; halosubstituted  $\text{C}_{1-6}$  alkyl, aryl, aryl $\text{C}_{1-6}$ alkyl, heteroaryl and heteroaryl $\text{C}_{1-6}$ alkyl; or

5 (vi) a bicyclic aryl, bicyclic heteroaryl, monocyclic or bicyclic reduced heteroaryl, or monocyclic heteroaryl group other than imidazolyl, optionally substituted by a group selected from  $\text{OR}^{17}$ ,  $-\text{COR}^{13}$ ,  $-\text{CO}_2\text{R}^{13}$ ,  $\text{SOR}^{13}$ ,  $\text{SO}_2\text{R}^{13}$ ,  $\text{CONR}^{13}\text{R}^{14}$ ,  $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$ ; halosubstituted  $\text{C}_{1-6}$  alkyl, aryl, aryl $\text{C}_{1-6}$ alkyl, heteroaryl and heteroaryl $\text{C}_{1-6}$ alkyl;

10

where  $\text{R}^{13}$  and  $\text{R}^{14}$  each independently represents hydrogen,  $\text{C}_{1-10}$ alkyl,  $\text{C}_{2-10}$ alkenyl,  $\text{C}_{2-10}$ alkynyl,  $\text{C}_{3-6}$ cycloalkyl,  $\text{C}_{3-6}$ cycloalkenyl, aryl, aryl $\text{C}_{1-10}$ alkyl, heteroaryl, heteroaryl $\text{C}_{1-10}$ alkyl, reduced heteroaryl or reduced heteroaryl,  $\text{C}_{1-10}$ alkyl, and  $\text{R}^{17}$  represents hydrogen or  $\text{C}_{2-10}$ alkenyl,

15  $\text{C}_{2-10}$ alkynyl,  $\text{C}_{3-6}$ cycloalkyl,  $\text{C}_{3-6}$ cycloalkenyl, aryl, aryl $\text{C}_{1-10}$ alkyl, heteroaryl, heteroaryl $\text{C}_{1-10}$ alkyl, reduced heteroaryl or reduced heteroaryl,  $\text{C}_{1-10}$ alkyl

and  $\text{R}^{8a}$ ,  $\text{R}^{9a}$ ,  $\text{R}^{10a}$  and  $\text{R}^{11a}$  are each independently hydrogen, halo, hydroxy, amino, nitro, cyano, thiol,  $\text{C}_{1-10}$ alkyl,  $\text{C}_{1-10}$ alkoxy,  $\text{C}_{1-10}$ cycloalkyl,  $\text{C}_{1-10}$ cycloalkoxy,  $\text{C}(\text{O})\text{R}^{15}$ ,

20  $\text{C}(\text{O})\text{NR}^{15}\text{R}^{16}$ ,  $\text{S}(\text{O})\text{R}^{15}$  or halo $\text{C}_{1-10}$ alkyl;

where  $\text{R}^{15}$  and  $\text{R}^{16}$  each independently represent hydrogen or  $\text{C}_{1-10}$ alkyl.

In compounds of formula (IIa):

25

When  $\text{R}^{1a}$  represents a group defined in (i) this is preferably a  $\text{C}_{10-25}$  e.g. a  $\text{C}_{10-20}$  branched or unbranched alkyl group, advantageously a  $\text{C}_{12-18}$  e.g. a  $\text{C}_{14-18}$  branched or unbranched alkyl group, optionally interrupted by one or more oxygen atoms.

Optional substituents for said alkyl groups are preferably selected from one or more of

30 aryl e.g. phenyl; heteroaryl e.g. thienyl; aryloxy, e.g. phenoxy; heteroaryloxy, e.g.

benzoxazinyloxy;  $-\text{CO}_2\text{R}^{13}$  e.g.  $\text{COOH}$ ;  $\text{NR}^{13}\text{R}^{14}$  e.g.  $\text{NH}_2$ ;  $-\text{CONR}^{13}\text{R}^{14}$  e.g.  $\text{CONHCH}_3$ ; cyano, nitro, halogen and OH.  $\text{R}^{13}$  and  $\text{R}^{14}$  preferably each independently represent hydrogen or  $\text{C}_{1-6}$ alkyl.

5      When  $\text{R}^{1a}$  represents a group defined by (ii) this is preferably a  $\text{C}_{10-25}$  e.g. a  $\text{C}_{10-20}$  branched or unbranched alkenyl or alkynyl group, advantageously a  $\text{C}_{14-18}$  branched or unbranched alkenyl or alkynyl group. Preferred optional substituents are those defined as preferred substituents in group (i).

10     When  $\text{R}^{1a}$  represents a group defined by (iii) the  $\text{C}_{1-9}$  group preferably contains one or two oxygen atoms. Preferred optional substituents are as defined above for groups (i) and (ii).

15     When  $\text{R}^{1a}$  represents a group defined by (iv) preferred substituents are as defined above for groups (i) - (iii).

20     When  $\text{R}^{1a}$  represents a group defined by (v) preferred substituents are selected from  $\text{OR}^{17}$ ,  $\text{CO}_2\text{R}^{13}$ ,  $\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$  and aryl  $\text{C}_{1-10}$ alkyl. The phenyl group  $\text{R}^{1a}$  and any additional phenyl group or moiety in the substituent may also be substituted by one or more of halogen, alkyl or haloalkyl.

When  $\text{R}^{1a}$  represents a group defined by (vi) this is preferably selected from naphthyl, pyridyl, pyrrolyl and piperidiny.

25      $\text{R}^{1a}$  is preferably selected from groups (i) and (v) defined above.

$\text{R}^{1a}$  is preferably  $\text{C}_{10-20}$  branched or unbranched alkyl, optionally interrupted by one or two oxygen atoms

and/or optionally substituted by one or more of aryl, eg. phenyl; aryloxy e.g. phenoxy wherein the phenyl moiety may be optionally substituted by alkyl, haloalkyl, halogen or phenoxy; heteroaryl, eg. thienyl; heteroaryloxy e.g. benzoxazinyloxy (which may optionally carry an oxo substituent); cyano, nitro,  $-\text{CO}_2\text{R}^{13}$ ,  $-\text{NR}^{13}\text{R}^{14}$ ,  $-\text{CONR}^{13}\text{R}^{14}$ ,  
5 OH and halogen.

$\text{R}^{1a}$  is also advantageously phenyl substituted by one or more, but most preferably one, of  $\text{OR}^{17}$ ,  $-\text{CO}_2\text{R}^{13}$ ,  $-\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$  and aryl  $\text{C}_{1-10}$ alkyl.

10  $\text{R}^{13}$  and  $\text{R}^{14}$  preferably each independently represent hydrogen or  $\text{C}_{1-6}$ alkyl.

$\text{R}^{17}$  preferably represents phenyl, optionally substituted by alkyl, haloalkyl, halogen or phenoxy, wherein the phenyl moiety may also be optionally substituted by alkyl, haloalkyl or halogen.

15

Most preferably  $\text{R}^{1a}$  is an unbranched alkyl chain having 14, 15, 16, 17 or 18 carbon atoms.

$\text{R}^{8a}$  is preferably hydrogen or halogen eg. fluorine; most preferably hydrogen.

20

$\text{R}^{9a}$  is preferably hydrogen or lower branched or unbranched alkyl having 1 to 10 carbon atoms, preferably 1 to 6 carbon atoms, preferably methyl, cyclic alkyl having 3 to 6 carbon atoms, preferably cyclopropyl, halo $\text{C}_{1-6}$ alkyl, preferably trifluoromethyl or a halogen, preferably chlorine or fluorine;

25

$\text{R}^{10a}$  is preferably hydrogen, lower branched or unbranched alkyl having 1 to 10 carbon atoms, preferably 1-6 carbon atoms preferably methyl, cyclic alkyl having 3 to 6 carbon atoms preferably cyclopropyl, halo $\text{C}_{1-6}$ alkyl preferably trifluoromethyl or a halogen preferably chlorine or fluorine;

30



$R^{11a}$  is preferably hydrogen, halogen, eg. fluorine; or branched or unbranched alkyl having 1 to 10 carbon atoms, preferably 1-6 carbon atoms, eg. methyl.

Preferably, in compounds of formula (IIa) at least one of  $R^{8a}$ ,  $R^{9a}$ ,  $R^{10a}$  and  $R^{11a}$  represents a substituent other than hydrogen. Thus, for example,  $R^{8a}$  may represent a hydrogen atom and  $R^{9a}$ ,  $R^{10a}$  and  $R^{11a}$  are as defined above. In a further preferred embodiment each of  $R^{8a}$  and  $R^{11a}$  represents a hydrogen atom,  $R^{9a}$  represents a substituent as defined above and  $R^{10a}$  represents a hydrogen atom or a substituent. In a further embodiment  $R^{8a}$ ,  $R^{9a}$  and  $R^{10a}$  represent hydrogen and  $R^{11a}$  represents a substituent as defined above, eg. methyl. Most preferably each of  $R^{8a}$ ,  $R^{10a}$  and  $R^{11a}$  represents a hydrogen atom, and  $R^{9a}$  represents a substituent as defined above, advantageously a  $C_{1-6}$ alkyl group, especially methyl.

A further group of novel compounds within the scope of formula (II) is that wherein  $R^1$  is as defined for formula (II) and at least one of  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  represents a  $C_{8-20}$  alkyl group, preferably a  $C_{8-10}$  alkyl group. Most preferably in this embodiment either  $R^9$  or  $R^{10}$  represents a  $C_{8-10}$  alkyl group and the remaining substituents on the benzene ring are all hydrogen.

Examples of pharmaceutically acceptable salts of the above compounds include those derived from organic acids such as methanesulphonic acid, benzenesulphonic acid and p-toluenesulphonic acid, mineral acids such as hydrochloric and sulphuric acid and the like, giving methanesulphonate, benzenesulphonate, p-toluenesulphonate, hydrochloride and sulphate, and the like, respectively or those derived from bases such as organic and inorganic bases. Examples of suitable inorganic bases for the formation of salts of compounds for this invention include the hydroxides, carbonates, and bicarbonates of ammonia, lithium, sodium, calcium, potassium, aluminium, iron, magnesium, zinc and the like. Salts can also be formed with suitable organic bases. Such bases suitable for the formation of pharmaceutically acceptable base addition salts with compounds of the present invention include organic bases which are








nontoxic and strong enough to form salts. Such organic bases are already well known in the art and may include amino acids such as arginine and lysine, mono-, di-, or trihydroxyalkylamines such as mono-, di-, and triethanolamine, choline, mono-, di-, and trialkylamines, such as methylamine, dimethylamine, and trimethylamine, 5 guanidine; N-methylglucosamine; N-methylpiperazine; morpholine; ethylenediamine; N-benzylphenethylamine; tris(hydroxymethyl) aminomethane; and the like.









Salts may be prepared in a conventional manner using methods well known in the art. Acid addition salts of said basic compounds may be prepared by dissolving the free 10 base compounds according to the first or second aspects of the invention in aqueous or aqueous alcohol solution or other suitable solvents containing the required acid. Where a compound of formula (I) contains an acidic function a base salt of said compound may be prepared by reacting said compound with a suitable base. The acid or base salt may separate directly or can be obtained by concentrating the solution eg. by 15 evaporation. The compounds of this invention may also exist in solvated or hydrated forms.










The invention also extends to prodrugs of the aforementioned compounds. A prodrug is commonly described as an inactive or protected derivative of an active ingredient or 20 a drug which is converted to the active ingredient or drug in the body.












Representative compounds according to the first and/or second aspects of the invention are those which include:










Table 1






Reference Number	Structure	Compound Name
1		2-Ethoxy-6-methyl-4 <i>H</i> -3,1-benzoxazin-4-one
2		2-Phenoxy-4 <i>H</i> -3,1-benzoxazin-4-one
3		2-(4-Methoxyphenoxy)-4 <i>H</i> -3,1-benzoxazin-4-one
4		2-(4-Methylphenoxy)-4 <i>H</i> -3,1-benzoxazin-4-one
5		2-(4-Chlorophenoxy)-4 <i>H</i> -3,1-benzoxazin-4-one
6		2-(2-Chloroethoxy)-4 <i>H</i> -3,1-benzoxazin-4-one
7		2-Propoxy-4 <i>H</i> -3,1-benzoxazin-4-one

8		6-Methyl-2-phenoxy-4 <i>H</i> -3,1-benzoxazin-4-one
9		6-Methyl-2-propoxy-4 <i>H</i> -3,1-benzoxazin-4-one
10		2-(2-Ethylhexyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
11		6-Methyl-2-octyloxy-4 <i>H</i> -3,1-benzoxazin-4-one
12		2-Hexyloxy-6-methyl-4 <i>H</i> -3,1-benzoxazin-4-one
13		2(2-Ethylhexyloxy)-6-methyl-4 <i>H</i> -3,1-benzoxazin-4-one
14		6-Ethyl-2-hexyloxy-4 <i>H</i> -3,1-benzoxazin-4-one
15		2-Decyloxy-6-methyl-4 <i>H</i> -3,1-benzoxazin-4-one









16		6-Methyl-2-tetradecyloxy-4H-3,1-benzoxazin-4-one
17		6-Methyl-2-pentadecyloxy-4H-3,1-benzoxazin-4-one
18		2-Hexadecyloxy-6-methyl-4H-3,1-benzoxazin-4-one
19		2-Heptadecyloxy-6-methyl-4H-3,1-benzoxazin-4-one
20		6-Methyl-2-octadecyloxy-4H-3,1-benzoxazin-4-one
21		7-Ethyl-2-hexyloxy-4H-3,1-benzoxazin-4-one
22		2-(3,7-Dimethyloctyloxy)-6-methyl-4H-3,1-benzoxazin-4-one
23		2-[2-(2-Hexyloxyethoxy)ethoxy]-6-methyl-4H-3,1-benzoxazin-4-one
24		(Z)-6-Methyl-2-(octadeca-9-enyloxy)-4H-3,1-benzoxazin-4-one









25		6-Methyl-2-(10-phenyldecyloxy)-4H-3,1-benzoxazin-4-one
26		7-Ethyl-2-octyloxy-4H-3,1-benzoxazin-4-one
27		2-Octyloxy-4H-3,1-benzoxazin-4-one
28		6-Methoxy-2-octyloxy-4H-3,1-benzoxazin-4-one
29		6-Methyl-2-(4-phenoxyphenoxy)-4H-3,1-benzoxazin-4-one
30		2-Hexyloxy-4H-3,1-benzoxazin-4-one
31		2-Dodecyloxy-6-methyl-4H-3,1-benzoxazin-4-one
32		6-Iodo-2-octyloxy-4H-3,1-benzoxazin-4-one
33		7-Butyl-2-Octyloxy-4H-3,1-benzoxazin-4-one
34		6-Methyl-2-(8-phenyloctyloxy)-4H-3,1-benzoxazin-4-one
35		6-Methyl-2-(4-phenylbutyloxy)-4H-3,1-benzoxazin-4-one








36		6-Methyl-2-(12-phenyldodecyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
37		( <i>Z</i> )-6-Methyl-2-(octadeca-11-enyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
38		6-Methyl-2-(octadeca-11-ynyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
39		6-Methyl-2-[10-(thien-2-yl)decyloxy]-4 <i>H</i> -3,1-benzoxazin-4-one
40		5-Fluoro-2-hexadecyloxy-4 <i>H</i> -3,1-benzoxazin-4-one
41		8-Fluoro-2-hexadecyloxy-4 <i>H</i> -3,1-benzoxazin-4-one
42		6-Fluoro-2-hexadecyloxy-4 <i>H</i> -3,1-benzoxazin-4-one
43		6-Chloro-2-hexadecyloxy-4 <i>H</i> -3,1-benzoxazin-4-one
44		6-Cyclopropyl-2-hexadecyloxy-4 <i>H</i> -3,1-benzoxazin-4-one








45		2-Hexadecyloxy-6-hydroxy-4H-3,1-benzoxazin-4-one
46		2-Hexadecyloxy-6-mercapto-4H-3,1-benzoxazin-4-one
47		6-Amino-2-hexadecyloxy-4H-3,1-benzoxazin-4-one
48		2-Hexadecyloxy-6-nitro-4H-3,1-benzoxazin-4-one
49		6-Cyano-2-hexadecyloxy-4H-3,1-benzoxazin-4-one






50		2-Hexadecyloxy-6-trifluoromethyl-4H-3,1-benzoxazin-4-one
51		6-Formyl-2-hexadecyloxy-4H-3,1-benzoxazin-4-one
52		6-Acetamido-2-hexadecyloxy-4H-3,1-benzoxazin-4-one
53		2-Hexadecyloxy-6-sulfo-4H-3,1-benzoxazin-4-one
54		2-Hexadecyloxy-7-trifluoromethyl-4H-3,1-benzoxazin-4-one
55		2-Hexadecyloxy-7-hydroxy-4H-3,1-benzoxazin-4-one
56		7-Amino-2-hexadecyloxy-4H-3,1-benzoxazin-4-one
57		7-Cyclopropyl-2-hexadecyloxy-4H-3,1-benzoxazin-4-one

58		7-Chloro-2-hexadecyloxy-4 <i>H</i> -3,1-benzoxazin-4-one
59		2-Hexadecyloxy-4 <i>H</i> -pyrido[2,3-d][1,3]oxazin-4-one
60		( <i>E</i> )-2-(Hexadeca-5-enyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
62		2-(2-Naphthyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
64		2-(3-Pyridyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
65		2-(2-Pyrrolyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
66		2-(2-Piperidinyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
67		2-[6-(2-Pyrrol)ylhexyloxy]-4 <i>H</i> -3,1-benzoxazin-4-one

69		2-(14-Cyanotetradecyloxy)-4H-3,1-benzoxazin-4-one
70		2-(14-Nitrotetradecyloxy)-4H-3,1-benzoxazin-4-one
71		2-(15-Methoxypentadecyloxy)-4H-3,1-benzoxazin-4-one
72		2-(15-Phenylpentadecyloxy)-4H-3,1-benzoxazin-4-one
73		2-(14-Carboxytetradecyloxy)-4H-3,1-benzoxazin-4-one
74		2-(14-Aminotetradecyloxy)-4H-3,1-benzoxazin-4-one
75		2-(14-Hydroxytetradecyloxy)-4H-3,1-benzoxazin-4-one

76		2-(12-N-Methylcarbamoyldodecyloxy)-4H-3,1-benzoxazin-4-one
77		2-Hexadecyloxy-6,7-dimethyl-4H-3,1-benzoxazin-4-one
78		5-Methyl-2-octyloxy-4H-3,1-benzoxazin-4-one
79		7-Octyl-2-octyloxy-4H-3,1-benzoxazin-4-one
80		6-Octyl-2-octyloxy-4H-3,1-benzoxazin-4-one
81		2-(5-Chloropentyloxy)-6-methyl-4H-3,1-benzoxazin-4-one
82		2,2'-(1,16-Hexadecylidenedioxy)-bis-4H-3,1-benzoxazin-4-one

83		6,8-Dimethyl-2-octyloxy-4 <i>H</i> -3,1-benzoxazin-4-one
84		6-Methyl-2-(6-phenoxyhexyloxy)-4 <i>H</i> -3,1-benzoxazin-4-one
85		6-Methyl-2-[6-(4-phenoxyphenoxy)hexyloxy]-4 <i>H</i> -3,1-benzoxazin-4-one

Apart from compounds 1-3, 6, 7 and 9, the compounds of Table 1 are believed to be novel and as such form preferred embodiments of the present invention.

Preferred compounds of formula (II) listed in Table 1 include compounds numbers 4,  
5 5, 8, 10, 11, 12, 13, 14, 21, 26, 27, 28, 30, 32, 33, 34, 35 and 78.

These specific compounds are believed to be novel and as such form a further aspect of the invention.

10 Preferred compounds of formula (IIa) listed in Table 1 include compounds numbers 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, 29, 31, 36, 37, 38, 39, 79, 80, 81, 82, 84 and 85.

Particularly preferred compounds of formula (IIa) are:

- 15 15: 2-Decyloxy-6-methyl-4*H*-3,1-benzoxazin-4-one  
16: 6-Methyl-2-tetradecyloxy-4*H*-3,1-benzoxazin-4-one  
18: 2-Hexadecyloxy-6-methyl-4*H*-3,1-benzoxazin-4-one


Preferred compounds of the invention listed above extend to the tautomers thereof, as  
20 well as (but not limited to) pharmaceutically acceptable salts, esters, amides or prodrugs thereof or a derivative optionally with one or more lipid groups (natural or synthetic) attached.

A third aspect of the invention provides a process for the manufacture of any one or  
25 more of the novel compounds or derivatives according to the first or second aspects of the invention. Thus, the present invention provides a process for the preparation of a novel compound of formula (II) in particular a compound of formula (IIa) which process comprises:

30 Process (A) reacting a compound of formula (IV):

29

5



(IV)

with a compound of formula (V):

10



(V)


15

or

Process (B) cyclising a compound of formula (VI)

20

25



(VI)


30

wherein R<sup>1</sup> and R<sup>8</sup>-R<sup>11</sup> are as hereinbefore defined and R<sup>18</sup> is hydrogen or C<sub>1-6</sub>alkyl.

or:

Process (C) reacting a compound of formula (VII)

5



10

with a compound of formula (VIII):



15

or:

Process (D) converting a compound of formula (I), (II) or (IIa) into a different compound of formula (IIa), by, for example,

20

(i) reduction of a compound of formula (IIa) wherein any of  $R^1$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  contains an alkenyl or alkynyl group or moiety, to the corresponding alkyl or alkenyl group or moiety; or

(ii) alkylation of a compound of formula (IIa) where one or more of  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  represents a halogen atom.

25

Process (A) may be effected by reacting a compound of formula (IV) with a chloroformate of formula (V). The process is preferably carried out under basic conditions, e.g. using pyridine. The reaction may be cooled to avoid overheating. The resulting carbamate intermediate is then cyclised by reaction with excess  
30 chloroformate or by addition of another cyclisation reagent, which promotes ring



closure. Suitable cyclisation reagents include for example, methyl chloroformate, carbonyl diimidazole, acetic anhydride, phosgene, oxalyl chloride, thionyl chloride or a peptide coupling agent such as dicyclohexyl carbodiimide (DCC). The cyclisation reagent is preferably phosgene, triphosgene or thionyl chloride. It will be understood  
5 by those skilled in the art that when a chloroformate is used to effect the cyclisation, this may be provided by employing an excess of the compound of formula (V). Preferably, however, a low molecular weight chloroformate is employed, on grounds of cost and ease of removing the resulting alcohol.

10 Compounds of formula (V) for use in the process (A) may be prepared by standard methods well known in the art, e.g. by reaction of the corresponding alcohol  $R^1OH$  with phosgene in a solvent such as toluene. The product may be isolated in conventional manner by removal of solvent and volatile by-products.

15 Process (B) may be effected by reaction of a compound (VI) wherein  $R^{18}$  is hydrogen, in the presence of a cyclisation reagent, e.g. an alkyl chloroformate, for example as described for process (A). Alternatively a compound (VI) may be cyclised by treatment with a dehydrating agent such as concentrated sulphuric acid.

20 Compounds (VI) wherein  $R^{18}$  is an alkyl group may be prepared by reacting an ester corresponding to formula (IV) with e.g. phosgene and a base such as pyridine to afford the corresponding isocyanate, followed by treatment with an alcohol  $R^1OH$ . If desired the ester (i.e. where  $R^{18}$  is alkyl) may be hydrolysed to the corresponding acid ( $R^{18}=H$ ) using for example lithium hydroxide in e.g. aqueous tetrahydrofuran or aqueous  
25 dioxane.


It will be appreciated that process (A) also proceeds via an intermediate of formula (VI) and is hence a variant of process (B).

Process (C) may be effected by reacting a compound of formula (VII) with an alcohol  $R^1OH$  in the presence of a base, e.g. Hunig's base (diisopropylethylamine).

A compound of formula (VII) may be prepared from the corresponding anhydride of formula (IX)

5

10



by reaction with for example phosphorus oxychloride ( $POCl_3$ ) at elevated temperature e.g.  $100^\circ C$ .

15

The anhydride of formula (IX) may itself be obtained by cyclisation of a compound of formula (IV), for example using phosgene or a synthetic equivalent.

20

In process (D), reduction of an alkenyl or alkynyl group may be effected for example by catalytic hydrogenation using e.g. 10% palladium on charcoal in an alcoholic solvent, such as ethanol, under 1 atmosphere of hydrogen gas.

25

Alkylation according to process (D)(ii) may be effected using a Stille or other palladium catalysed cross-coupling process, using e.g. tetra-alkyl tin such as tetramethyl tin and  $PhCH_2Pd(PPh_3)_2Cl$  in HMPA at elevated temperature e.g.  $50-100^\circ C$ . Other halides or pseudohalides e.g. triflates may be employed as starting materials.

30

A fourth aspect of the invention is a compound according to the first and second aspects of the invention (i.e. compounds of formulae (I), (II) and (IIa)), for use in

medicine. Preferred features of the first and second aspects of the invention also apply to the fourth aspect. Further details of the fourth aspect of the invention are set out in the text which follows.

5 A fifth aspect of the invention relates to a compound according to the first and/or second aspects of the invention for use in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality. This includes both *in vivo* and *in vitro* uses and other uses such as industrial uses. Such an enzyme is one which catalyses the breakdown of a substrate containing an ester functionality by  
10 the addition of water, resulting in the cleavage of a chemical bond. Such enzymes are involved in key processes in the body. Enzymes according to this invention include lipases (hydrolyse fatty acid esters), esterases (hydrolyse esters) and phosphatases (hydrolyse phosphate esters).

15 The enzyme is preferably a lipase. Lipases include pancreatic lipase, gastric lipase, lipoprotein lipase, lingual lipase, adipose tissue lipase, hormone sensitive lipase, phospholipase A1, A2, B, C, D etc., hepatic lipase, and other triacyl, diacyl and monoacylglycerol lipases in the mammalian body. Many similar such lipases are also known in plants, fungi and microorganisms.

20 Also covered are esterase enzymes and phosphatase enzymes. Esterase enzymes include pig liver esterase, cholesteryl esterase, retinyl esterase, 1-alkyl-2-acetylgllycerophosphocholine esterase, carboxylic ester hydrolases, and cholesterol esterase. Phosphatase enzymes include serine/threonine phosphatases PP1, PP2 and  
25 PP3, phosphoprotein phosphatase, myosin-light-chain phosphatase, protein phosphoprotein 2C, and protein tyrosine phosphatase.

Compounds according to the invention, for use in medicine, are primarily for use in relation to the prevention and/or treatment of a medical condition such as obesity,  
30 hyperlipaemia, hyperlipidaemia and related diseases such as hyperglycaemia (type II

diabetes), hypertension, cardiovascular disease, stroke, gastrointestinal disease and gastrointestinal conditions. Compounds according to the first and second aspect of the invention are useful in these and other conditions due to their ability to inhibit an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester  
5 functionality (*in vivo*, as the enzyme naturally occurs). The invention also relates to non-medical weight loss, such as cosmetic weight loss and includes improving bodily appearance in general. Throughout this text, the prevention and/or treatment of any disorder means any effect which mitigates any damage or any medical disorder, to any extent, and includes prevention and treatment themselves. The term "treatment"  
10 means any amelioration of disorder, disease, syndrome, condition, pain or a combination of two or more thereof.

Clearly, an important application of the invention is in relation to weight loss (of all kinds as described above) in humans. However, the invention applies to medical and  
15 non-medical weight loss in any animal whose metabolism of fat and fat derivatives involves an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality (*in vivo*, as the enzyme naturally occurs). Thus, the invention has veterinary application and is particularly useful in relation to medical and non-medical weight loss in companion animals such as pet cats and dogs as well as in animals  
20 which provide meat for human consumption. In the case of the latter, the application of the present invention is to reduce fat content in order to provide a leaner meat product.

It is also believed that the compounds may be useful in reducing levels of toxins (e.g.  
25 dioxins and PCBs) stored in body fat. Without wishing to be bound by theory, it is believed that increasing the amount of undigested fat passing through the body enhances diffusion of toxins from fat stored in the body into fats in the blood, and thence into the intestine.

The fifth aspect of the invention has important applications. It includes test and diagnostic methods and the control and inhibition of unwanted enzymes, preferably lipases, in any process or in any product. The processes or products, which preferably involve a lipase, include: processing of agricultural commodities (e.g. oilseeds),  
5 recovery and isolation of enzymes from biotechnological processes (e.g. involving lysis of microorganisms), the manufacture and extraction of crude oil (especially oil and plastics), the industrial manufacture of triglycerides or other fats, manufacture of healthcare goods which comprise surfactants, soap or detergent (e.g. bath oils, creams), the manufacturing and processing of liposomes (e.g. healthcare products,  
10 diagnostics, gene therapy), the treatment of industrial waste (e.g. paper effluent treatment) and preventing the degradation of foodstuff which comprises a fat (e.g. chocolate processing). Thus, the invention also relates to these products and processes, e.g. a foodstuff which comprises a compound according to the first aspect of the invention, in particular foodstuffs which have a high fat content such as cakes,  
15 biscuits, pastry-products and the like and chocolate products. The preferred features of the fifth aspect of the invention, including an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality (*in vivo*, as the enzyme naturally occurs) are as discussed for the previous aspects of the invention.

20 A sixth aspect of the invention provides a composition comprising a novel compound according to the first or second aspect of the invention, in combination with a pharmaceutically acceptable carrier or diluent. Suitable carriers and/or diluents are well known in the art and include pharmaceutical grade starch, mannitol, lactose, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, (or other  
25 sugar), magnesium carbonate, gelatin, oil, alcohol, detergents, emulsifiers or water (preferably sterile). The composition may be a mixed preparation of a composition or may be a combined preparation for simultaneous, separate or sequential use (including administration).

The compounds according to the invention for use in the aforementioned indications may be administered by any convenient method, for example by oral (including by inhalation), parenteral, mucosal (e.g. buccal, sublingual, nasal), rectal or transdermal administration and the compositions adapted accordingly.

5

For oral administration, the compounds can be formulated as liquids or solids, for example solutions, syrups, suspensions or emulsions, tablets, capsules and lozenges.

10

A liquid formulation will generally consist of a suspension or solution of the compound or physiologically acceptable salt in a suitable aqueous or non-aqueous liquid carrier(s) for example water, ethanol, glycerine, polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

15

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and microcrystalline cellulose.

20

A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, powders, granules or pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

25

Compositions for oral administration may be designed to protect the active ingredient against degradation as it passes through the alimentary tract, for example by an outer coating of the formulation on a tablet or capsule.

30

Typical parenteral compositions consist of a solution or suspension of the compound or physiologically acceptable salt in a sterile aqueous or non-aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

Compositions for nasal or oral administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a pharmaceutically acceptable propellant. The aerosol dosage forms can also take the form of a pump-atomiser.

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

Compositions for rectal or vaginal administration are conveniently in the form of suppositories (containing a conventional suppository base such as cocoa butter), pessaries, vaginal tabs, foams or enemas.

Compositions suitable for transdermal administration include ointments, gels, patches and injections including powder injections.

Conveniently the composition is in unit dose form such as a tablet, capsule or ampoule.

5 The compositions of the sixth aspect of the invention are useful in the prevention and/or treatment of obesity, obesity-related disorder, other medical weight loss and non-medical related weight loss. Preferred features of this aspect of the invention are as described above for the first to fifth aspects of the invention.

10 A seventh aspect of the invention provides a process for the manufacture of a composition according to the sixth aspect of the invention. The manufacture can be carried out by standard techniques well known in the art and involves combining a compound according to the first or second aspect of the invention and the pharmaceutically acceptable carrier or diluent. The composition may be in any form including a tablet, a liquid, a capsule, and a powder or in the form of a food product,  
15 e.g. a functional food. In the latter case the food product itself may act as the pharmaceutically acceptable carrier.

An eighth aspect of the invention provides a method for the prevention and/or treatment of obesity or an obesity-related disorder, the method comprising the  
20 administration of a compound according to the first or second aspect of the invention, preferably in combination with a pharmaceutically acceptable carrier or diluent (as per the sixth aspect of the invention). Obesity-related disorders include hyperlipemia, hyperlipideamia, hyperglycaemia, hypertension, cardiovascular disease, stroke, gastrointestinal disease and gastrointestinal conditions. The compound or composition  
25 is preferably administered to a patient in need thereof and in a quantity sufficient to prevent and/or treat the symptoms of the condition, disorder or disease. For all aspects of the invention, particularly medical ones, the administration of a compound or composition has a dosage regime which will ultimately be determined by the attending physician and will take into consideration such factors such as the compound being  
30 used, animal type, age, weight, severity of symptoms, method of administration,



adverse reactions and/or other contraindications. Specific defined dosage ranges can be determined by standard design clinical trials with patient progress and recovery being fully monitored. Such trials may use an escalating dose design using a low percentage of the maximum tolerated dose in animals as the starting dose in man.

5

The physiologically acceptable compounds of the invention will normally be administered in a daily dosage regimen (for an adult patient) of, for example, an oral dose of between 1 mg and 2000 mg, preferably between 30 mg and 1000 mg, e.g. between 10 and 250 mg or an intravenous, subcutaneous, or intramuscular dose of  
10 between 0.1 mg and 100 mg, preferably between 0.1 mg and 50 mg, e.g. between 1 and 25 mg of the compound of the formula (I) or a physiologically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

15

A ninth aspect of the invention provides a cosmetic method (non-therapeutic) for maintaining a given weight, or for cosmetic weight loss, the method comprising the administration of a compound according to the first or second aspect of the invention, preferably in combination with a pharmaceutically acceptable carrier or diluent (as per  
20 the sixth aspect of the invention). The compound or composition is preferably administered to a subject in need or in desideratum thereof and in a quantity sufficient to maintain a given weight or for cosmetic weight loss.

25

The eighth and ninth aspects of the invention relate to methods which are applicable to humans and other animals, in particular companion animals (such as dogs and cats) and other animals which provide meat for human consumption, such as cattle, pigs and sheep (all of any age).

30

The invention will now be described with reference to the following non-limiting examples.

## Biological Test Methods and Results

### Test Compounds

5 The benzoxazinone compounds used in the following tests are identified by the reference number assigned in Table 1 hereinbefore.

### Measurement of lipase activity using a quinine diimine dye colorimetric assay

10 The inhibitory activity of the selected compounds to pancreatic lipase was measured in the following assay available from Sigma Ltd (Lipase-PS™, catalog number 805-A):

## Pancreatic lipase

$$\text{1,2-dibutyryn} \text{-----} > \text{2-monglyceride} + \text{fatty acid}$$

### Monoglyceride lipase

$$\text{2-monoglyceride} \xrightarrow{\text{hydrolysis}} \text{glycerol} + \text{fatty acid}$$

## Glycerol kinase

20      glycerol + ATP -----> glycerol-3-phosphate + ADP

### Glycerol phosphate oxidase

$$\text{glycerol-3-phosphate} + \text{O}_2 \text{ ----- } > \text{dihydroxyacetone phosphate} + \text{H}_2\text{O}_2$$

## Peroxidase

$$\text{H}_2\text{O}_2 + 4\text{-AAP} + \text{TOOS} \longrightarrow \text{quinine diimine dye} + 4\text{H}_2\text{O}$$

The glycerol released from the action of pancreatic and monoglyceride lipase was oxidised to release  $\text{H}_2\text{O}_2$ . The peroxidase reaction step then produces a quinine dye which is pink in colour and absorbs light at a wavelength of 550 nm.

### Inhibitor

Individual test compounds were dissolved in DMSO (dimethyl sulphoxide) at 10 mM. DMSO was used to avoid any problems with compounds being water-insoluble.

- 5 For individual compounds, the  $IC_{50}$  (concentration at which lipase activity is inhibited to one half of the maximum) was calculated by measuring the inhibitory activity from log-dose response curves using a range of inhibitor concentrations.

### Results

10

A range of compounds was assayed in the quinine diimine dye colorimetric assay which provides a rapid method to measure lipase inhibitory activity. None of the compounds tested interfered with the colorimetric reaction, i.e. they did not give false positive results.

15

A range of inhibitory activities for the tested benzoxazinone compounds was observed, indicating that these compounds are inhibitors of human pancreatic lipase. The following compounds had an  $IC_{50}$  of  $\leq 100$  nM:

Compounds 9, 11-16, 18-22, 23, 24-39, 77, 78, 79-82, 84 and 85.

20

### **Measurement of lipase enzyme activity using a NaOH titration method**

The inhibitory activity of the selected compounds to pancreatic lipase was measured in the assay described in Pasquier *et al* ; 1996, Vol 7, *Nutritional Biochemistry*, 293-302.

25

Log dose/response curves were constructed using a range of inhibitor concentrations.

## Results

Selected benzoxazinone compounds were tested in the NaOH titration assay. In this assay, the activity of porcine pancreatic lipase in a system containing lipid micelles is recorded. These conditions are therefore similar to those encountered in the gastrointestinal tract.

A range of inhibitory activities were observed for the tested benzoxazinone compounds in this assay, indicating that these compounds are inhibitors of porcine pancreatic lipase. The following compounds had  $IC_{50}$ 's of  $\leq 1 \mu M$ :

Compounds 1, 2, 4, 6-9, 11-16, 18, 22-25, 27-36, 37-39, 78, 82, 84 and 85.

The results demonstrate that a number of selected benzoxazinones are inhibitors of fat digestion and that these compounds may be particularly suitable for the treatment of obesity.

### *Measurement of trypsin activity*

Porcine trypsin (Boehringer) was dissolved at a concentration of 1 mg/ml in 100 mM MOPS (3-[N-Morpholino]propanesulphonic acid) pH 7.3 containing 2 mM  $CaCl_2$ . Prior to use, the enzyme was diluted 500 times to give a final concentration of 2  $\mu g/ml$ .

Selected compounds were routinely stored as 5 mM stock solutions dissolved in DMSO (Dimethylsulphoxide) at  $-20^\circ C$ . For the assay, aliquots were defrosted and a series of dilutions (x100, x200, x1,000, x2,000, x10,000, x20,000 and x100,000) made in 100 mM MOPS pH 7.3 containing 2 mM  $CaCl_2$ . The substrate Bz-Phe-Val-Arg-pNA (Benzoyl-phenylalanyl-valyl-arginine-*p*-nitroanilide) was dissolved in DMSO to

give a 10 mM solution. Immediately prior to use, the substrate was diluted to 0.3mM (30  $\mu$ l/ml) in 100 mM MOPS containing 2 M  $\text{CaCl}_2$ .

The assay was set up in triplicate in a 96 well ELISA plate. 10  $\mu$ l 2 $\mu$ g/ml trypsin, 26  $\mu$ l diluted inhibitor and 190  $\mu$ l substrate were added sequentially. The plates were then incubated at 37 °C in a BioRad Benchmark Microplate Reader. The rate of release of *p*-nitroaniline was measured at 405 nM over 10 minutes relative to that of the enzyme without inhibitor.

#### 10 *Measurement of chymotrypsin activity*

Bovine Chymotrypsin (Sigma Type 11 Cat. No. C4129) was dissolved at a concentration of 1 mg/ml in 100mM Tris pH 7.8. Prior to use, the enzyme was diluted 20 fold immediately before use, with the same buffer.

15

Selected compounds were routinely stored as 5 mM stock solutions dissolved in DMSO (Dimethylsulphoxide) at -20°C. For the assay, aliquots were defrosted and a series of dilutions (x20, x100, x200, x1,000, x2,000, x10,000, x20,000 and x100,000) made in 100 mM Tris pH 7.8. The substrate H-Ala-Ala-Phe-*p*-nitroanilide (H-alanyl-alanyl-phenylalanine-*p*-nitroanilide) (Bachem Cat. No. L-1095) was dissolved in DMSO to give a 10mM stock and stored at 4°C until use. Immediately prior to use, the substrate was diluted to 0.3mM final concentration (30  $\mu$ l/ml) just before use.

20

The assay was set up in triplicate in a 96 well ELISA plate. 10  $\mu$ l 50 $\mu$ g/ml chymotrypsin, 50  $\mu$ l diluted inhibitor and 190  $\mu$ l substrate were added sequentially. The plates were then incubated at 37 °C in a BioRad Benchmark Microplate Reader. The rate of release of *p*-nitroaniline was measured at 405 nM over 10 minutes relative to that of the enzyme without inhibitor.

25

## Results

The selectivity of compound 18 towards human pancreatic lipase, trypsin and chymotrypsin was measured in the assays described above. The inhibitory activity of compound 18 towards lipase was assessed in the quinine diimine dye colorimetric assay.

At 500 nM compound 18 had inhibited 98.7% of the pancreatic lipase activity but only 4% of the trypsin activity and 12.5% of the chymotrypsin activity. Thus compound 18 is a highly selective lipase inhibitor.

## Mouse Model Assay

Compound 18 was assayed in a mouse model as described by Isler *et al.*, *British Journal of Nutrition*, 1995, 73:851-862 and was found to be a potent lipase inhibitor.

## Synthesis of Intermediates

### Synthesis of 4-substituted anthranilic acids

*Example: 4-octyl anthranilic acid (4-octyl-2-aminobenzoic acid)*

Method based on that of L.A. Paquette et al. *J.Am.Chem.Soc.* 99, 3734 (1981)



A solution of 1-bromo-4-octylbenzene (9.9g, 36mmol) in sulfuric acid (20ml) was cooled in an ice bath. To this was added nitric acid (1.44ml, 36mmol). The ice bath was removed and the mixture stirred at room temperature for 20 minutes. A further

45

portion of nitric acid was added (0.07ml, 1.75mmol), stirring being continued for a further 20 min. The mixture was poured into aqueous potassium carbonate, which was extracted with ethyl acetate. The organic extract was washed with saturated aqueous potassium carbonate, water and brine then dried ( $\text{MgSO}_4$ ) and concentrated.


- 5 Purification of the crude product by flash chromatography (1% EtOAc/hexane) removed the unwanted (major) regioisomer and afforded the desired material as a yellow oil (1.7g, 5.4mmol).



10

The substrate (1.7g, 5.4mmol), copper (I) cyanide (0.533g, 5.9mmol) and pyridine (20ml) were refluxed at 150°C for 2 days. Concentration *in vacuo* and purification by flash chromatography (10% to 20% EtOAc/hexane) gave the desired material as a brown oil (739mg, 2.8mmol)

15



20

The substrate (694mg, 2.7mmol) was heated at 150°C in a mixture of water (2ml), AcOH (1ml) and sulfuric acid (1ml) for 2 days. The mixture was extracted with ethyl acetate, the organic phase being washed with water (x 2), dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated to give the desired material (744mg, 2.7mmol).

25

46




The starting material (744mg, 2.7mmol) was dissolved in ethanol (10ml) and to this  
 5 was added a slurry of 10% palladium on charcoal (40mg) in ethanol (4ml). The flask  
 was flushed with nitrogen then hydrogen (1 atm) after which stirring was maintained  
 overnight. Further portions of catalyst (5mg and 25mg) were added, the reaction being  
 complete after a further 24h. The reaction mixture was filtered through celite,  
 thoroughly rinsing with methanol and ethyl acetate. Concentration gave the anthranilic  
 10 acid (597mg, 2.4mmol) of sufficient purity for use without further purification;  $\delta_H$   
 (400 MHz,  $CDCl_3$ ) 0.79-0.81 (3H, m, Me), 1.12-1.36 (10H, m, 5 x  $CH_2$ ), 1.52 (2H,  
 br.s,  $ArCH_2CH_2$ ), 2.45 (2H, br.s,  $ArCH_2$ ), 6.42 (2H, br.s., 2 x  $ArH$ ), 7.74 (1H, br.s,  
 $ArH$ );  $m/z$  ( $ES^+$ ) 250 ( $MH^+$ ).

### 15 Synthesis of 5-substituted anthranilic acids

#### *Example: 5-octyl anthranilic acid*

Method based on that of B.R. Baker et al. J.Org.Chem 17, 141 (1952)




Chloral hydrate (3.97g, 24mmol) was dissolved in water (50ml). To this solution was  
 added, sequentially, anhydrous sodium sulfate (5.5g, 39mmol), 4-octylaniline (5ml,  
 22mmol), water (15ml), concentrated hydrochloric acid (2.3ml) and an aqueous  
 25 solution of hydroxylamine hydrochloride (4.5g in 22ml, 65mmol). The heterogeneous  
 mixture was heated to 95°C for 2 h, then 110°C for a further 1 h. The reaction mixture



47

was cooled to room temperature, the brown precipitate filtered and washed with water. This was dissolved in dichloromethane, dried ( $\text{MgSO}_4$ ) and concentrated to give 5.6g of crude material which was purified by flash chromatography on silica (20% EtOAc/hexane) to afford the desired material (2g, 7.2mmol).


5



10

The oxime (1.8g, 6.5mmol) was added to a mixture of conc. sulfuric acid (13ml) and water (1ml) at  $60^\circ\text{C}$  over the course of 15 min. The mixture was then heated to  $80^\circ\text{C}$  for 2h then stood at room temperature overnight. This was then extracted with ethyl acetate (x3), the combined organic layers being washed with saturated aqueous sodium bicarbonate and water until the washings were neutral. The organic phase was dried ( $\text{MgSO}_4$ ) and concentrated to give the isatin as a red solid (1.5g, 5.8mmol), which was used without further purification.

15



20

A mixture of the isatin (1.5g, 5.8mmol) and 1.5M sodium hydroxide (13ml) was warmed to  $50^\circ\text{C}$ . The heat was removed and a solution of 35% aqueous hydrogen peroxide (1.5ml) was added, at a rate that maintained the temperature at  $50$ - $55^\circ\text{C}$ . The reaction was then allowed to cool and stirred at room temperature for 30 min.


25

Acidification to pH 2 with concentrated hydrochloric acid caused the product to precipitate. The liquid was decanted off, the solid being washed with water. The solid

was partitioned between water and dichloromethane, the organic phase being washed with brine, dried (MgSO<sub>4</sub>) and concentrated to give the desired anthranilic acid (1.4g, 5.6mmol). No further purification was required:  $\delta$  (400 MHz, CDCl<sub>3</sub>) 0.81 (3H, t, *J* 6.6, Me), 1.20-1.23 (10H, m, 5 x CH<sub>2</sub>), 1.49 (2H, br.s, ArCH<sub>2</sub>CH<sub>2</sub>), 2.41-2.44 (2H, m, ArCH<sub>2</sub>), 6.55 (1H, d, *J* 8.3, ArH), 7.09 (1H, d, *J* 8.3, ArH), 7.65 (1H, s, ArH); *m/z* (ES<sup>+</sup>) 250 (MH<sup>+</sup>).

Preparation of aryl chloroformates:


Example: 4-Phenoxyphenylchloroformate



4-Phenoxyphenol (1.68g, 9 mmol), 1,4-dimethylimidazolidin-2-one (0.051 ml, 0.45 mmol) and phosgene solution (4.5 ml of a 20% solution in toluene, 9 mmol) were heated to 40 C for 30 min. The temperature was then increased to 80 C, and five further portions of phosgene solution (2.25 ml each, 4.5 mmol) were added at 30 min intervals. 30 min after the last addition, the solution was allowed to cool to r.t. and stand overnight. An aliquot was added to MeOH/pyridine to form the stable methyl carbamate, whereupon tlc (10% EtOAc/hexane) showed almost complete disappearance of starting material. The chloroformate solution was used directly in the preparation of compound 29, using the procedure described in Example 4 below.


Synthesis of novel compounds according to the invention

Example 1

6-Methyl-2-octyloxy-4H-3,1-benzoxazin-4-one (reference number 11)

A solution of 2-amino-5-methylbenzoic acid (302 mg, 2 mmol) in pyridine (10 ml) was cooled to 0°C and treated dropwise with octyl chloroformate (1.15 ml, 6 mmol). The resulting mixture was allowed to warm to room temperature and stirred for 4 h. The pyridine was removed under vacuum and the residue dissolved in ethyl acetate (50 ml). This solution was washed with 1M HCl (10 ml) and brine (5 ml), dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford a pale orange oily solid.

Recrystallisation from hexane afforded the desired product as an off-white solid (144 mg, 25%);  $\delta_H$  (400 MHz, DMSO-d<sub>6</sub>) 0.68 (3H, t, *J* 7, CH<sub>2</sub>CH<sub>3</sub>), 1.26-1.40 (10H, m, 5 x CH<sub>2</sub>), 1.73 (2H, tt, *J*, *J'* 7, OCH<sub>2</sub>CH<sub>2</sub>), 2.35 (3H, s, CH<sub>3</sub>), 4.35 (2H, t, *J* 7, OCH<sub>2</sub>), 7.34 (1H, d, *J* 8, Ph), 7.65 (1H, d, *J* 8, Ph), 7.83 (1H, s, Ph); *m/z* (ES<sup>+</sup>) 290 (MH<sup>+</sup>).

Example 26-Methyl-2-phenoxy-4H-3,1-benzoxazin-4-one (reference number 8)

A solution of 2-amino-5-methylbenzoic acid (1.0 g, 6.6 mmol) in pyridine (10 ml) was cooled to 0°C and treated dropwise with phenyl chloroformate (3.3 ml, 26 mmol). The resulting mixture was allowed to warm to room temperature and stirred for 16 h, after which time the pyridine was removed *in vacuo*. The residue was washed with water (20 ml) and dried *in vacuo*. Recrystallisation from toluene afforded the desired product as a pale brown solid (692 mg, 41%);  $\delta_H$  (400 MHz, DMSO-d<sub>6</sub>) 2.40 (3H, s, CH<sub>3</sub>),


50

7.33-7.45 (3H, m, Ph), 7.48-7.55 (3H, m, Ph), 7.63 (1H, d,  $J$  8, Ph), 7.89 (1H, s, Ph- $H_5$ );  $m/z$  ( $ES^+$ ) 254 ( $MH^+$ ).

### Example 3

5

2-Propoxy-6-methyl-4*H*-3,1-benzoxazin-4-one (reference number 9)



10 A solution of 2-amino-5-methylbenzoic acid (1.0 g, 6.6 mmol) in pyridine (10 ml) was cooled to 0°C and treated dropwise with propyl chloroformate (3.0 ml, 26 mmol). The resulting mixture was allowed to warm to room temperature and stirred for 4 h, after which time the pyridine was removed *in vacuo*. The residue was washed with water (25 ml) and dried *in vacuo* to afford the desired product as an off-white solid (0.96 g, 15 66%);  $\delta_H$  (400 MHz, DMSO- $d_6$ ) 1.03 (3H, t,  $J$  7,  $CH_2CH_3$ ), 1.82 (2H, tq,  $J$ ,  $J'$  7,  $CH_2CH_3$ ), 2.46 (3H, s,  $CH_3$ ), 4.42 (2H, t,  $J$  7,  $OCH_2$ ), 7.40 (1H, d,  $J$  8, Ph), 7.71 (1H, d,  $J$  8, Ph), 7.89 (1H, s, Ph);  $m/z$  ( $ES^+$ ) 219 ( $MH^+$ ).


### Example 4

20

2-Hexadecyloxy-6-methyl-4*H*-3,1-benzoxazin-4-one (Compound 18)

### Preparation 1


Step 1:



25

1-Hexadecanol (0.78 g, 3.2 mmol, 1 eq.) was dissolved in the minimum amount of THF under nitrogen. To this was added a 20% solution of phosgene in toluene (2.34 ml, 5 mmol, 1.5 eq.). After 45 minutes, a second similar portion of phosgene solution was added. After a further 45 minutes, the apparatus was purged with nitrogen  
 5 (scrubbed on exit with 5M sodium hydroxide) to remove excess phosgene.

Step 2:




2-Amino-5-methylbenzoic acid (100 mg, 0.64 mmol, 0.2-eq.) was dissolved in pyridine (10 ml). The chloroformate solution was added dropwise by syringe and the mixture stirred at room temperature for 2 hours. The mixture was diluted with ethyl acetate (100 ml) and washed with 10% aqueous citric acid (100 ml, x2), saturated  
 15 sodium bicarbonate solution (100 ml, x2), water (100 ml) and saturated brine (100 ml). The organic phase was dried ( $\text{MgSO}_4$ ) and concentrated in vacuo.

The residue was purified by flash chromatography on silica, eluting with 1:5:94 diisopropylethylamine/ethyl acetate/hexane to afford a white solid (40mg, 15%)  $\delta_{\text{H}}$   
 20 (400 MHz,  $\text{CDCl}_3$ ) 0.87 (3H, t, J 6.8,  $\text{CH}_2\text{CH}_3$ ), 1.24-1.45 (26H, m,  $13 \times \text{CH}_2$ ), 1.75-1.83 (2H, m,  $\text{OCH}_2\text{CH}_2$ ), 2.41 (3H, s,  $\text{ArCH}_3$ ), 4.41 (2H, t, J 6.7,  $\text{OCH}_2$ ), 7.30 (1H, d, J 8.3, ArH), 7.51 (1H, dd, J 8.5, 2.0, ArH), 7.90 (1H, d, J 1.1, ArH);  $m/z$  ( $\text{ES}^+$ ) 402 ( $\text{MH}^+$ ); MPt. 72-73 °C.

Thin layer chromatograms (solvent 1% diisopropylamine/ 5% ethyl acetate/ 94% hexane) were visualised with UV and phosphomolybdic acid in ethanol ( $R_f$   
 25 (compound 18) = 0.6).

Preparation 2

Step 1:




5

1-Hexadecanol (5.01 g, 20.6 mmol, 1 eq.) was dissolved in THF (10 ml) under nitrogen and added to a 20% solution of phosgene in toluene (29 ml, 62.5 mmol, 3 eq.). The mixture was stirred at room temperature for 2 hours, then the apparatus was purged with nitrogen (scrubbed on exit with 5M sodium hydroxide) to remove excess phosgene.

10

Step 2:



2-Amino-5-methylbenzoic acid (2.71 g, 17.9 mmol, 0.87 eq.) was dissolved in pyridine (24 ml) and added to the chloroformate solution prepared above. The mixture was stirred at room temperature for 1.75 hours. Methyl chloroformate (13.6 ml, 176 mmol, 8.5 eq) was added slowly, then the mixture was left to stir at ambient temperature overnight. The mixture was diluted with ethyl acetate (20 ml) and washed with water (15 ml) and 10% aqueous citric acid (20 ml). The combined aqueous phases were extracted with ethyl acetate (20 ml). The organic extracts were combined and washed with water (20 ml) and brine (20 ml) then concentrated to give a solid. This was slurried in pentane (5 ml), filtered, then slurried in acetonitrile (5 ml), filtered and purified by flash chromatography on silica (1.5% diisopropylethylamine in dichloromethane) to give a white solid (2.51 g, 31%)  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 0.87 (3H, t, J 6.8,  $\text{CH}_2\text{CH}_3$ ), 1.24-1.45 (26H, m,  $13 \times \text{CH}_2$ ), 1.75-1.83 (2H, m,  $\text{OCH}_2\text{CH}_2$ ), 2.41

20

25

(3H, s, ArCH<sub>3</sub>), 4.41 (2H, t, J 6.7, OCH<sub>2</sub>), 7.30 (1H, d, J 8.3, ArH), 7.51 (1H, dd, J 8.5, 2.0, ArH), 7.90 (1H, d, J 1.1, ArH); m/z (ES<sup>+</sup>) 402 (MH<sup>+</sup>); MPt. 72-73 °C.

### Example 5

5

The other compounds listed in Table 1 may be prepared in a similar manner to Examples 1 to 4 above, in particular the following compounds were prepared using the starting materials indicated:

Compound number	Starting material 1	Starting material 2
4	2-aminobenzoic acid	4-methylphenyl chloroformate
5	2-aminobenzoic acid	4-chlorophenyl chloroformate
8	2-amino-5-methylbenzoic acid	phenyl chloroformate
10	2-aminobenzoic acid	2-ethylhexyl chloroformate
11	2-amino-5-methylbenzoic acid	octyl chloroformate
12	2-amino-5-methylbenzoic acid	hexyl chloroformate
13	2-amino-5-methylbenzoic acid	2-ethylhexyl chloroformate
14	2-amino-5-ethylbenzoic acid	hexyl chloroformate
15	2-amino-5-methylbenzoic acid	Decanol
16	2-amino-5-methylbenzoic acid	1-tetradecanol
17	2-amino-5-methylbenzoic acid	1-pentadecanol
19	2-amino-5-methylbenzoic acid	1-heptadecanol
20	2-amino-5-methylbenzoic acid	1-octadecanol
21	4-ethyl-2-aminobenzoic acid	Hexyl chloroformate
22	5-methyl-2-aminobenzoic acid	3,7-dimethyloctan-1-ol
23	5-methyl-2-aminobenzoic acid	2-(2-hexyloxyethoxy)ethanol
24	5-methyl-2-aminobenzoic acid	Z-9-octadecen-1-ol
25	5-methyl-2-aminobenzoic acid	10-phenyldecan-1-ol
26	4-ethyl-2-aminobenzoic acid	Octyl chloroformate


27	2-aminobenzoic acid	Octyl chloroformate
28	5-methoxy-2-aminobenzoic acid	Octyl chloroformate
29	5-methyl-2-aminobenzoic acid	4-phenoxyphenol
30	2-aminobenzoic acid	Hexyl chloroformate
31	2-aminobenzoic acid	1-dodecanol
32	5-iodo-2-aminobenzoic acid	Octyl chloroformate
33	4-butyl-2-aminobenzoic acid	Octyl chloroformate
34	5-methyl-2-aminobenzoic acid	8-phenyloctan-1-ol
35	5-methyl-2-aminobenzoic acid	4-phenylbutan-1-ol
36	5-methyl-2-aminobenzoic acid	12-phenyldodecan-1-ol
37	5-methyl-2-aminobenzoic acid	Z-11-octadecen-1-ol
38	5-methyl-2-aminobenzoic acid	11-octadecyn-1-ol
39	5-methyl-2-aminobenzoic acid	10-(2-thienyl)decan-1-ol
78	6-methyl-2-aminobenzoic acid	Octyl chloroformate
79	4-octyl-2-aminobenzoic acid	Octyl chloroformate
80	5-octyl-2-aminobenzoic acid	Octyl chloroformate
81	5-methyl-2-aminobenzoic acid	5-chloropentan-1-ol
82	2-amino-5-methylbenzoic acid	1,16-hexadecandiol
83	3,5-dimethyl-2-aminobenzoic acid	Octyl chloroformate
84	2-amino-5-methylbenzoic acid	6-phenoxyhexan-1-ol
85	2-amino-5-methylbenzoic acid	6-(4-phenoxyphenoxy)hexan-1-ol

The foregoing description details specific compounds, compositions, methods and uses which can be employed to practice the present invention. However, those skilled in the art will know how to use alternative reliable methods for aiming at alternative embodiments of the invention which are herein encompassed.



Claims

1. The use of a compound comprising formula (I):



5

(I)

- or a pharmaceutically acceptable salt, ester, amide or prodrug thereof; in the manufacture of a medicament for the treatment of conditions which require the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality

10

wherein in formula (I):

- 15 A is a 6-membered aromatic or heteroaromatic ring; and

- R<sup>1</sup> is a branched or unbranched alkyl (optionally interrupted by one or more oxygen atoms), alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, reduced arylalkyl, arylalkenyl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, reduced aryl, reduced heteroaryl, reduced heteroarylalkyl or a substituted derivative of any of the foregoing groups, wherein the substituents are one or more independently of halogen, alkyl, halosubstituted alkyl, aryl, arylalkyl, heteroaryl, reduced heteroaryl, reduced heteroarylalkyl, arylalkoxy, cyano, nitro, -C(O)R<sup>4</sup>, -CO<sub>2</sub>R<sup>5</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -NR<sup>6</sup>R<sup>7</sup>, -OR<sup>6</sup>, -SR<sup>6</sup>, -C(O)CX<sup>1</sup>X<sup>2</sup>NR<sup>6</sup>R<sup>7</sup>, -C(O)N(OH)R<sup>6</sup>, -C(O)NR<sup>5</sup>R<sup>4</sup>, -NR<sup>6</sup>C(O)R<sup>4</sup>, -CR<sup>6</sup>(NH<sub>2</sub>)CO<sub>2</sub>R<sup>6</sup>, -NHCX<sup>1</sup>X<sup>2</sup>CO<sub>2</sub>R<sup>6</sup>, -N(OH)C(O)NR<sup>6</sup>R<sup>7</sup>, -N(OH)C(O)R<sup>4</sup>, -NHC(O)NR<sup>6</sup>R<sup>7</sup>, -C(O)NHNHNR<sup>6</sup>R<sup>7</sup>, -C(O)N(OR<sup>5</sup>)R<sup>6</sup>, or a lipid or steroid (natural or

20

25

synthetic) with the proviso that any hetero atom substituent in  $R^1$  must be separated from the exocyclic oxygen atom by at least two carbon atoms (preferably saturated);

and where:-

5

$R^4$  is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, reduced heteroaryl, reduced heteroarylalkyl,  $-OR^6$ ,  $-NHCX^1X^2CO_2R^6$  or  $-NR^6R^7$ ;

10

$R^5$  is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, reduced heteroaryl or reduced heteroarylalkyl; and

15


$R^6$  and  $R^7$  are independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, reduced heteroaryl, reduced heteroarylalkyl or  $-(CH_2)_n(OR^5)_m$  wherein  $n$  is 1 to 12, preferably 2 to 10, wherein  $m$  is 1-3 and  $R^5$  is most preferably  $C_2-C_{10}$  alkyl; and

20

$X^1$  and  $X^2$  are independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, reduced heteroaryl or reduced heteroarylalkyl.

2. Use according to claim 1 wherein the compound (I) is a compound of formula (II):

25



(II)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof,

5

wherein:-

$R^1$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$ ,  $X^1$  and  $X^2$  are as defined above for formula (I); and

10  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{11}$  are each independently hydrogen, halo, hydroxy, amino, nitro, cyano,

or a group  $R^1$ , as defined above,

or a group  $R^{12}Q$  where  $Q$  is O, CO, CONH, NHCO, S, SO,  $SO_2$ , or  $SO_2NH_2$  and  $R^{12}$  is

15 hydrogen or a group  $R^1$  as defined above,

or a group  $R^1R^2N$  where  $R^1$  is as defined above and  $R^2$  is hydrogen or  $R^1$ , with the proviso that any hetero atom substituent in  $R^1$  and/or  $R^2$  must be separated from the aromatic hetero atom substituent by at least two carbon atoms (preferably saturated).

20

3. Use according to claim 2, wherein in the compound of formula (II),  $R^1$  is either a branched or unbranched alkyl group having up to 25 carbon atoms; an aryl group; an arylalkyl group wherein the alkyl moiety has up to 25 carbon atoms; or an aryl aryl group, wherein the aryl alkyl group or the aryl aryl group may be separated by a

spacer, and where the spacer can be one or more of an ester, amide, O, CH<sub>2</sub> or a ketone;

R<sup>8</sup> is hydrogen or fluorine;

5

R<sup>9</sup> is lower branched or unbranched alkyl having 1 to 10 carbon atoms; cyclic alkyl having 3 to 10 carbon atoms; haloalkyl; or a halogen;

10

R<sup>10</sup> is hydrogen lower branched or unbranched alkyl having 1 to 10 carbon atoms; cyclic alkyl having 3 to 10 carbon atoms; haloalkyl; or a halogen; and

R<sup>11</sup> is hydrogen lower branched or unbranched alkyl having 1 to 10 carbon atoms or a halogen.

15

4. Use according to any of claims 1 to 3 wherein said condition is selected from obesity, hyperlipaemia, hyperlipidaemia, hyperglycaemia (type II diabetes), hypertension, cardiovascular disease, stroke, gastrointestinal disease and gastrointestinal conditions.

20

5. Use according to any of claims 1 to 3 wherein said medicament is for reducing levels of toxins in body fat.


6. Use according to any of claims 1 to 5 wherein the medicament is for administration to humans.

25

7. Use according to any of claims 1 to 5 wherein the medicament is for administration to animals.

8. A compound formula (IIa):

30



(IIa)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof; where:

5  $R^{1a}$  is

- (ii) a  $C_{10-30}$  branched or unbranched alkyl, optionally substituted by one or more independently of  $C_{3-6}$  cycloalkyl,  $C_{3-6}$  cycloalkenyl, aryl, heteroaryl, reduced heteroaryl,  $-C(O)R^{13}$ ,  $-CO_2R^{13}$ ,  $-SOR^{13}$ ,  $-SO_2R^{13}$ ,  $-NR^{13}R^{14}$ ,  $-OR^{13}$ ,  $-SR^{13}$ ,  $-C(O)NR^{13}R^{14}$ ,  $-NR^{14}C(O)R^{13}$ , halogen, cyano, and nitro and/or optionally interrupted by one or more oxygen atoms with the proviso that any hetero atom in  $R^1$  must be separated from the exocyclic oxygen atom (or from any other heteroatom) by at least two carbon atoms (preferably saturated);
- 10
- (ii)  $C_{2-25}$  alkenyl,  $C_{2-25}$  alkynyl,  $C_{3-6}$  cycloalkenyl, aryl- $C_{2-25}$  alkenyl, heteroaryl- $C_{2-25}$  alkenyl, reduced heteroaryl, reduced heteroaryl- $C_{1-25}$  alkyl or a substituted derivative of any of the foregoing groups wherein the substituents are one or more independently of  $C_{1-6}$  alkyl, halosubstituted  $C_{1-6}$  alkyl, aryl, aryl- $C_{1-6}$  alkyl, heteroaryl, reduced heteroaryl, reduced heteroaryl- $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, aryl- $C_{1-6}$  alkoxy,  $-C(O)R^{13}$ ,  $-CO_2R^{13}$ ,  $-SOR^{13}$ ,  $-SO_2R^{13}$ ,  $-NR^{13}R^{14}$ ,  $-OR^{13}$ ,  $-SR^{13}$ ,  $-C(O)NR^{13}R^{14}$ ,  $-NR^{14}C(O)R^{13}$ , halogen, cyano, and nitro, with the proviso that any hetero atom in  $R^1$  must be separated from the exocyclic oxygen atom (or from any other heteroatom) by at least two carbon atoms (preferably saturated);
- 15
- 20

- (iii) a  $C_{1-9}$  alkyl group interrupted by one or more oxygen atoms and optionally substituted by one or more independently of  $C_{3-6}$  cycloalkyl,  $C_{3-6}$  cycloalkenyl, aryl, heteroaryl, reduced heteroaryl,  $-C(O)R^{13}$ ,  $-CO_2R^{13}$ ,  $-SOR^{13}$ ,  $-SO_2R^{13}$ ,  $NR^{13}R^{14}$ ,  $OR^{13}$ ,  $SR^{13}$ ,  $-C(O)NR^{13}R^{14}$ ,  $-NR^{14}C(O)R^{13}$ , halogen, cyano and nitro with the proviso that any hetero atom in  $R^1$  must be separated from the exocyclic oxygen atom (or from any other heteroatom) by at least two carbon atoms (preferably saturated); or
- (iv) a  $C_{1-9}$  alkyl group substituted by a group selected from  $-C(O)R^{13}$ ,  $-CO_2R^{13}$ ,  $SOR^{13}$ ,  $SO_2R^{13}$ ,  $NR^{13}R^{14}$ ,  $OR^{13}$ ,  $SR^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ; tetrahydronaphthyl, pyridyl, pyrrolyl, piperidiny, halogen, cyano, nitro, bicyclic aryl, bicyclic heteroaryl, monocyclic or bicyclic reduced heteroaryl, monocyclic heteroaryl other than imidazolyl;
- (v) a phenyl group substituted by a group selected from  $OR^{17}$ ,  $-COR^{13}$ ,  $-CO_2R^{13}$ ,  $SOR^{13}$ ,  $SO_2R^{13}$ ,  $CONR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ; halosubstituted  $C_{1-6}$  alkyl, aryl, aryl $C_{1-6}$ alkyl, heteroaryl and heteroaryl $C_{1-6}$ alkyl; or
- (vi) a bicyclic aryl, bicyclic heteroaryl, monocyclic or bicyclic reduced heteroaryl, or monocyclic heteroaryl group other than imidazolyl, optionally substituted by a group selected from  $OR^{17}$ ,  $-COR^{13}$ ,  $-CO_2R^{13}$ ,  $SOR^{13}$ ,  $SO_2R^{13}$ ,  $CONR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ; halosubstituted  $C_{1-6}$  alkyl, aryl, aryl $C_{1-6}$ alkyl, heteroaryl and heteroaryl $C_{1-6}$ alkyl;
- where  $R^{13}$  and  $R^{14}$  each independently represents hydrogen,  $C_{1-10}$ alkyl,  $C_{2-10}$ alkenyl,  $C_{2-10}$ alkynyl,  $C_{3-6}$ cycloalkyl,  $C_{3-6}$ cycloalkenyl, aryl, aryl $C_{1-10}$ alkyl, heteroaryl, heteroaryl $C_{1-10}$ alkyl, reduced heteroaryl or reduced heteroaryl,  $C_{1-10}$ alkyl, and  $R^{17}$  represents hydrogen or  $C_{2-10}$ alkenyl,

C<sub>2-10</sub>alkynyl, C<sub>3-6</sub>cycloalkyl, C<sub>3-6</sub>cycloalkenyl, aryl, arylC<sub>1-10</sub>alkyl, heteroaryl, heteroarylC<sub>1-10</sub>alkyl, reduced heteroaryl or reduced heteroaryl, C<sub>1-10</sub>alkyl

5 and R<sup>8a</sup>, R<sup>9a</sup>, R<sup>10a</sup> and R<sup>11a</sup> are each independently hydrogen, halo, hydroxy, amino, nitro, cyano, thiol, C<sub>1-10</sub>alkyl, C<sub>1-10</sub>alkoxy, C<sub>1-10</sub>cycloalkyl, C<sub>1-10</sub>cycloalkoxy, C(O)R<sup>15</sup>, C(O)NR<sup>15</sup>R<sup>16</sup>, S(O)R<sup>15</sup> or haloC<sub>1-10</sub>alkyl;

where R<sup>15</sup> and R<sup>16</sup> each independently represent hydrogen or C<sub>1-10</sub>alkyl.

10 9. A compound according to claim 8 wherein R<sup>1a</sup> is C<sub>10-20</sub> branched or unbranched alkyl, optionally interrupted by one or two oxygen atoms and/or optionally substituted by one or more of aryl, aryloxy, heteroaryl, heteroaryloxy, cyano, nitro, CO<sub>2</sub>R<sup>13</sup>, -NR<sup>13</sup>R<sup>14</sup>, -CONR<sup>13</sup>R<sup>14</sup>, OH and halogen, wherein R<sup>13</sup> and R<sup>14</sup> each independently represent hydrogen or C<sub>1-6</sub>alkyl.

15

10. A compound according to claim 8 or 9 wherein R<sup>1a</sup> is an unbranched alkyl chain having 14, 15, 16, 17 or 18 carbon atoms.

20

11. A compound according to any of claims 8 to 10 wherein R<sup>8a</sup> is hydrogen or halogen.

25


12. A compound according to any of claims 8 to 11 wherein R<sup>9a</sup> is hydrogen; lower branched or unbranched alkyl having 1 to 10 carbon atoms; cyclic alkyl having 3 to 6 carbon atoms; haloC<sub>1-6</sub>alkyl or a halogen.

13. A compound according to any of claims 8 to 12 wherein R<sup>10a</sup> is hydrogen; lower branched or unbranched alkyl having 1 to 10 carbon atoms; cyclic alkyl having 3 to 6 carbon atoms; haloC<sub>1-6</sub>alkyl or a halogen.

14. A compound according to any of claims 8 to 13 wherein  $R^{11a}$  is hydrogen, halogen or branched or unbranched alkyl having 1 to 10 carbon atoms.

15. A compound of formula (IIb)

5



10

wherein  $R^1$  is as defined for formula (II) and at least one of  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  represents a  $C_{8-20}$  alkyl group, preferably a  $C_{8-10}$  alkyl group.

16. A novel compound of formula (II), selected from:

15

2-(4-Methylphenoxy)-4*H*-3,1-benzoxazin-4-one;

2-(4-Chlorophenoxy)-4*H*-3,1-benzoxazin-4-one,

6-Methyl-2-phenoxy-4*H*-3,1-benzoxazin-4-one,

2-(2-Ethylhexyloxy)-4*H*-3,1-benzoxazin-4-one,

6-Methyl-2-octyloxy-4*H*-3,1-benzoxazin-4-one,

20

2-Hexyloxy-6-methyl-4*H*-3,1-benzoxazin-4-one,

2(2-Ethylhexyloxy)-6-methyl-4*H*-3,1-benzoxazin-4-one,

2-Ethyl-2-hexyloxy-4*H*-3,1-benzoxazin-4-one,

7-Ethyl-2-hexyloxy-4*H*-3,1-benzoxazin-4-one,

7-Ethyl-2-octyloxy-4*H*-3,1-benzoxazin-4-one,

25

2-Octyloxy-4*H*-3,1-benzoxazin-4-one,

6-Methoxy-2-octyloxy-4*H*-3,1-benzoxazin-4-one,

2-Hexyloxy-4*H*-3,1-benzoxazin-4-one,

6-Iodo-2-octyloxy-4*H*-3,1-benzoxazin-4-one,

2-Octyloxy-7-propyl-4*H*-3,1-benzoxazin-4-one,

30

6-Methyl-2-(8-phenyloctyloxy)-4*H*-3,1-benzoxazin-4-one,



6-Methyl-2-(4-phenylbutoxy)-4*H*-3,1-benzoxazin-4-one,  
5-Methyl-2-octyloxy-4*H*-3,1-benzoxazin-4-one;  
or a pharmaceutically acceptable salt, ester, amide or prodrug thereof.


- 5      17.    A compound of formula (IIa) selected from:
- 2-Decyloxy-6-methyl-4*H*-3,1-benzoxazin-4-one,  
6-Methyl-2-tetradecyloxy-4*H*-3,1-benzoxazin-4-one,  
6-Methyl-2-pentadecyloxy-4*H*-3,1-benzoxazin-4-one,  
2-Heptadecyloxy-6-methyl-4*H*-3,1-benzoxazin-4-one,  
10    6-Methyl-2-octadecyloxy-4*H*-3,1-benzoxazin-4-one,  
2-(3,7-Dimethyloctyloxy)-6-methyl-4*H*-3,1-benzoxazin-4-one,  
2-[2-(2-Hexyloxyethoxy)ethoxy]-6-methyl-4*H*-3,1-benzoxazin-4-one,  
2-(Octadeca-9-enyloxy)-6-methyl-4*H*-3,1-benzoxazin-4-one,  
2-(10-Phenyldecyloxy)-6-methyl-4*H*-3,1-benzoxazin-4-one,  
15    6-Methyl-2-(4-phenoxyphenoxy)-4*H*-3,1-benzoxazin-4-one,  
2-Dodecyloxy-6-methyl-4*H*-3,1-benzoxazin-4-one,  
6-Methyl-2-(12-phenyldodecyloxy)-4*H*-3,1-benzoxazin-4-one, 6-Methyl-2-(octadeca-  
11-enyloxy)-4*H*-3,1-benzoxazin-4-one,  
6-Methyl-2-(octadeca-11-ynyloxy)-4*H*-3,1-benzoxazin-4-one, 6-Methyl-2-[-10-(thien-  
20    2-yl)-decyloxy]-4*H*-3,1-benzoxazin-4-one,  
7-Octyl-2-octyloxy-4*H*-3,1-benzoxazin-4-one,  
6-Octyl-2-octyloxy-4*H*-3,1-benzoxazin-4-one,  
2-(5-Chloropentyloxy)-6-methyl-4*H*-3,1-benzoxazin-4-one,  
2,2'-(1,16-Hexadecylidenedioxy)-*bis*-4*H*-3,1-benzoxazin-4-one,  
25    6-Methyl-2-(6-phenoxyhexyloxy)-4*H*-3,1-benzoxazin-4-one,  
6-Methyl-2-[6-(4-phenoxyphenoxy)hexyloxy]-4*H*-3,1-benzoxazin-4-one;  
or a pharmaceutically acceptable salt, ester amide or prodrug thereof.

18. A compound of formula (IIa) which is:

2-Hexadecyloxy-6-methyl-4*H*-3,1-benzoxazin-4-one or a pharmaceutically acceptable salt, ester, amid or prodrug thereof.

5 19. A process for the preparation of a novel compound of formula (II) which process comprises:

Process (A) reacting a compound of formula (IV):




15 with a compound of formula (V):



20 or


Process (B). cyclising a compound of formula (VI)



30 wherein R<sup>1</sup> and R<sup>8</sup>-R<sup>11</sup> are as hereinbefore defined and R<sup>18</sup> is hydrogen or C<sub>1-6</sub>alkyl.

or:

Process (C) reacting a compound of formula (VII)



10 with a compound of (VIII)



15 or:

Process (D) converting a compound of formula (I), (II) or (IIa) into a different compound of formula (IIa)

20 20. A compound according to any of claims 8 to 18 , for use in medicine.

21. A pharmaceutical composition comprising a novel compound of formula (II) or a pharmaceutically acceptable salt, ester, amide or pro-drug thereof, in combination with a pharmaceutically acceptable carrier or diluent.

25 22. A food product comprising a compound of formula (II) or a pharmaceutically acceptable salt, ester, amide or pro-drug thereof.

23. A method for the prevention or treatment of obesity or an obesity related disorder, the method comprising administering a compound, as defined in any one of

30

claims 1 to 18, or a composition as claimed in claim 21 or claim 22 to a patient.

24. A compound, as defined in any one of claims 1 to 18, for use in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester  
5 functionality.

25. Use of a compound as defined in any of claims 1 to 18 or a pharmaceutically acceptable salt, ester, amide or prodrug thereof to reduce fat content of animals which provide meat for human consumption.

10

26. A cosmetic method for maintaining a given weight, or for cosmetic weight loss, the method comprising the administration of a compound as defined in any of claims 1 to 18.

15 27. A compound comprising formula (I) hereinbefore described with reference to one or more of the examples.

28. A process for obtaining a compound comprising formula (I) hereinbefore described with reference to one or more of the examples.

20

29. The use of a compound comprising formula (I) in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality hereinbefore described with reference to one or more of the examples.

25 30. Use according to claim 29 of a compound of formula (I) in the control and inhibition of unwated enzymes in a process or product.

31. Use according to claim 30 in the manufacture of healthcare goods comprising surfactants, soap or detergents.

30

32. Use according to claim 30 in preventing the degradation of foodstuff which comprises a fat.

33. A novel intermediate of formula (VI) as hereinbefore defined.

# INTERNATIONAL SEARCH REPORT

Inter.      nal Application No

PCT/GB 00/00031

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D265/26 C07C271/28 A61K31/536 A61P3/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	M. GÜTSCHOW ET AL.: "Inhibition of cathepsin G by 4H-3,1-benzoxazin-4-ones" BIOORGANIC & MEDICINAL CHEMISTRY, vol. 5, no. 10, 1997, pages 1935-1942, XP000904822 the whole document	1-33
X	R.L. JARVEST ET AL.: "Inhibition of HSV-1 protease by benzoxazinones" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 6, no. 20, 1996, pages 2463-2466, XP000904816 the whole document	1-33
A	DD 246 996 A (KARL MARX UNI) 24 June 1987 (1987-06-24) cited in the application the whole document	1-33
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 April 2000

Date of mailing of the international search report

28/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

# INTERNATIONAL SEARCH REPORT

Inter:      nal Application No

PCT/GB 00/00031

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 206 323 A (SYNTEX INC) 30 December 1986 (1986-12-30) claims & US 4 665 070 A cited in the application ---	1-33
P,A	WO 99 16758 A (REDDY S RESEARCH FOUNDATION DR ;REDDY CHEMINOR INC (US)) 8 April 1999 (1999-04-08) claims -----	1-33

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 00/00031

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 23,26  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 23,26  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☒ Claims Nos.: 28-32  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:  
Claims 28-32 were not searched since they refer to any compounds,  
formulations, processes and methods mentioned in the description;  
therefore it does not comply with the requirements of the PCT, Rule 6.2a.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 28-32

Claims 28-32 were not searched since they refer to any compounds, formulations, processes and methods mentioned in the description; therefore it does not comply with the requirements of the PCT, Rule 6.2a.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. Patent Application No

PCT/GB 00/00031

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DD 246996	A	24-06-1987	NONE	
EP 0206323	A	30-12-1986	US 4665070 A	12-05-1987
			AU 4912090 A	31-05-1990
			AU 599320 B	19-07-1990
			AU 5915086 A	08-01-1987
			CA 1270487 A	19-06-1990
			DK 297486 A	26-12-1986
			ES 556514 D	01-05-1988
			ES 8802223 A	01-07-1988
			ES 557730 D	01-03-1988
			ES 8801806 A	01-05-1988
			ES 557731 D	01-03-1988
			ES 8801807 A	01-05-1988
			ES 557732 D	01-03-1988
			ES 8801808 A	01-05-1988
			FI 862691 A	26-12-1986
			HU 41750 A,B	28-05-1987
			JP 62030770 A	09-02-1987
			KR 9004319 B	22-06-1990
			KR 9004803 B	06-07-1990
			NO 862534 A,B,	29-12-1986
			NO 871576 A,B,	29-12-1986
			NZ 230589 A	21-12-1989
			US 4745116 A	17-05-1988
			ZA 8604715 A	24-02-1988
WO 9916758	A	08-04-1999	NONE	

REVISED VERSION

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
13 July 2000 (13.07.2000)

PCT

(10) International Publication Number  
WO 00/40569 A1(51) International Patent Classification<sup>7</sup>: C07D 265/24,  
498/04, 413/12, A61K 31/536, A61P 3/04 // (C07D  
498/04, 265/00, 221:00)Therapeutics Limited, 280 Cambridge Science Park,  
Milton Road, Cambridge CB4 4WE (GB).

(21) International Application Number: PCT/GB00/00031

(74) Agents: FLORENCE, Julia, Anne et al.; Kilburn &  
Strode, 20 Red Lion Street, London WC1R 4PJ (GB).

(22) International Filing Date: 6 January 2000 (06.01.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
9900416.0 8 January 1999 (08.01.1999) GB(71) Applicant (for all designated States except US): AL-  
IZYME THERAPEUTICS LIMITED [GB/GB]; Ganta  
Park, Great Abingdon, Cambridge CB1 6GS (GB).

(72) Inventors; and


(75) Inventors/Applicants (for US only): HODSON, Harold,  
Francis [GB/GB]; 69 Whitecroft Way, Beckenham, Kent  
BR3 3AW (GB). DOWNHAM, Robert [GB/GB]; Cam-  
bridge Discovery Chemistry, The Merrifield Centre, Rose-  
mary Lane, Cambridge CB1 3LQ (GB). MITCHELL,  
Timothy, John [GB/GB]; Cambridge Discovery Chem-  
istry, The Merrifield Centre, Rosemary Lane, Cambridge  
CB1 3LQ (GB). CARR, Beverley, Jane [GB/GB];  
Boys Bridge Cottage, Fen Road, Abington Piggotts,  
Cambs. SG8 5PQ (GB). DUNK, Christopher, Robert  
[GB/GB]; Alizyme Therapeutics Limited, 280 Cambridge  
Science Park, Milton Road, Cambridge CB4 4WE (GB).  
PALMER, Richard, Michael, John [GB/GB]; Alizyme(81) Designated States (national): AL, AM, AT, AU, AZ, BA,  
BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES,  
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,  
SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

(88) Date of publication of the revised international search  
report: 15 March 2001(15) Information about Correction:  
see PCT Gazette No. 11/2001 of 15 March 2001, Section  
IIFor two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: 2-AMINO-BENZOXAZINONE DERIVATIVES FOR THE TREATMENT OF OBESITY

(57) Abstract: The use of a compound comprising formula (I) or a salt, ester, amide or prodrug thereof in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality, e.g. in the control and inhibition of unwanted enzymes in products and processes. The compounds are also useful in medicine e.g. in the treatment of obesity and related conditions. The invention also relates to novel compounds within formula (I), to processes for preparing them and pharmaceutical compositions containing them. In formula (I) A is a 6-membered aromatic or heteroaromatic ring; and R<sup>1</sup> is a branched or unbranched alkyl (optionally interrupted by one or more oxygen atoms), alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, reduced arylalkyl, arylalkenyl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, reduced aryl, reduced heteroaryl, reduced heteroarylalkyl or a substituted derivative of any of the foregoing groups.

WO 00/40569 A1

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC 7 C07D265/24 C07D498/04 C07D413/12 A61K31/536 A61P3/04  
 //(C07D498/04,265:00,221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GUETSCHOW M ET AL: "INHIBITION OF CATHEPSIN G BY 4H-3,1-BENZOXAZIN-4-ONES" BIOORGANIC & MEDICINAL CHEMISTRY,GB,ELSEVIER SCIENCE LTD, vol. 5, no. 10, 1997, pages 1935-1942, XP000904822 ISSN: 0968-0896 the whole document	1-32
X	JARVEST R L ET AL: "INHIBITION OF HSV-1 PROTEASE BY BENZOXAZINONES" BIOORGANIC & MEDICINAL CHEMISTRY,GB,ELSEVIER SCIENCE LTD, vol. 6, no. 20, 1996, pages 2463-2466, XP000904816 ISSN: 0968-0896 the whole document	1-32
	--- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

11 December 2000

Date of mailing of the international search report

12 12 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

## INTERNATIONAL SEARCH REPORT

In International Application No

PCT/GB 00/00031

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 37485 A (SEARLE & CO ;ABOOD NORMAN (US); FLYNN DANIEL L (US); BECKER DANIEL) 28 November 1996 (1996-11-28) claims ---	1-32
X	WO 96 07648 A (WARNER LAMBERT CO) 14 March 1996 (1996-03-14) claims & US 5 652 237 A 29 July 1997 (1997-07-29) cited in the application ---	1-32
X	EP 0 147 211 A (SYNTEX INC) 3 July 1985 (1985-07-03) claims & US 4 657 893 A 14 April 1987 (1987-04-14) cited in the application ---	1-32
X	DE 23 15 303 A (BAYER AG) 17 October 1974 (1974-10-17) cited in the application claims ---	1,19
X	US 3 450 700 A (ULRICH HENRI ET AL) 17 June 1969 (1969-06-17) the whole document -----	1-32

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/GB 00/00031

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 23,25,26,30-32 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 27-29  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 27-29

Claims 27-29 were not searched since they refer to any compounds, formulations, processes and methods mentioned in the description; therefore it does not comply with the requirements of the PCT, Rule 6.2a

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/GB 00/00031

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9637485	A	28-11-1996	US 5985872 A	16-11-1999
			AU 711174 B	07-10-1999
			AU 5803096 A	11-12-1996
			BR 9608506 A	15-06-1999
			CA 2222034 A	28-11-1996
			CN 1190963 A	19-08-1998
			EP 0828721 A	18-03-1998
			JP 11505849 T	25-05-1999
			NZ 308398 A	29-06-1999
WO 9607648	A	14-03-1996	US 5652237 A	29-07-1997
			AU 3496295 A	27-03-1996
EP 0147211	A	03-07-1985	US 4657893 A	14-04-1987
			AT 56444 T	15-09-1990
			AU 586616 B	20-07-1989
			AU 3716984 A	04-07-1985
			CA 1269800 A	29-05-1990
			DE 3483204 D	18-10-1990
			DK 625184 A	28-06-1985
			ES 539038 D	01-06-1986
			ES 8607260 A	01-11-1986
			ES 550879 D	01-03-1987
			ES 8703437 A	01-05-1987
			FI 845116 A,B,	28-06-1985
			HU 36808 A	28-10-1985
			HU 195648 B	28-06-1988
			IL 73943 A	31-01-1989
			JP 60169469 A	02-09-1985
			KR 8900992 B	15-04-1989
			NO 845176 A,C	28-06-1985
			NZ 210669 A	30-05-1988
			ZA 8410089 A	27-08-1986
DE 2315303	A	17-10-1974	NONE	
US 3450700	A	17-06-1969	NONE	